

Association Between Use of Enhanced Recovery After Surgery Protocol and Postoperative Complications in Total Hip and Knee Arthroplasty in the Postoperative Outcomes Within Enhanced Recovery After Surgery Protocol in Elective Total Hip and Knee Arthroplasty Study (POWER2)

Javier Ripollés-Melchor, MD; Ane Abad-Motos, MD; Yolanda Díez-Remesal, MD, PhD; Marta Aseguinolaza-Pagola, MD; Lidia Padin-Barreiro, MD; Rubén Sánchez-Martín, MD; Margarita Logroño-Egea, MD; Juan C. Catalá-Bauset, MD; Silvia García-Orallo, MD; Elvira Bisbe, MD, PhD; Nuria Martín, MD; Alejandro Suárez-de-la-Rica, PhD; Ana B. Cuéllar-Martínez, MD; Silvia Gil-Trujillo, MD; Juan Carlos Estupiñán-Jiménez, MD; Marta Villanova-Baraza, MD; Cristina Gil-Lapetra, MD; Pilar Pérez-Sánchez, MD; Nicolás Rodríguez-García, MD; Alvaro Ramiro-Ruiz, MD; Carla Farré-Tebar, MD; Alejandro Martínez-García, MD; Pedro Arauzo-Pérez, MD; Cristina García-Pérez, MD; Alfredo Abad-Gurumeta, MD, PhD; María A. Miñambres-Villar, MD; Alberto Sánchez-Campos, MD; Ignacio Jiménez-López, MD, PhD; José M. Tena-Guerrero, MD; Oliver Marín-Peña, MD, PhD; Miriam Sánchez-Merchante, MD; Ubaldo Vicente-Gutiérrez, MD; María C. Cassinello-Ogea, MD, PhD; Carlos Ferrando-Ortolá, MD, PhD; Héctor Berges-Gutiérrez, MD; Jesús Fernanz-Antón, MD; Manuel A. Gómez-Ríos, MD; Daniel Bordonaba-Bosque, MD, PhD; José M. Ramírez-Rodríguez, MD, PhD; José Antonio García-Erce, PhD; César Aldecoa, MD, PhD; for the Postoperative Outcomes Within Enhanced Recovery After Surgery Protocol in Elective Total Hip and Knee Arthroplasty (POWER2) Study Investigators Group for the Spanish Perioperative Audit and Research Network (REDGERM)

+ Supplemental content

IMPORTANCE The Enhanced Recovery After Surgery (ERAS) care protocol has been shown to improve outcomes compared with traditional care in certain types of surgery.

OBJECTIVE To assess the association of use of the ERAS protocols with complications in patients undergoing elective total hip arthroplasty (THA) and total knee arthroplasty (TKA).

DESIGN, SETTING, AND PARTICIPANTS This multicenter, prospective cohort study included patients recruited from 131 centers in Spain from October 22 through December 22, 2018. All consecutive adults scheduled for elective THA or TKA were eligible for inclusion. Patients were stratified between those treated in a self-designated ERAS center (ERAS group) and those treated in a non-ERAS center (non-ERAS group). Data were analyzed from June 15 through September 15, 2019.

EXPOSURES Total hip or knee arthroplasty and perioperative management. Sixteen individual ERAS items were assessed in all included patients, whether they were treated at a center that was part of an established ERAS protocol or not.

MAIN OUTCOMES AND MEASURES The primary outcome was postoperative complications within 30 days after surgery. Secondary outcomes included length of stay and mortality.

RESULTS During the 2-month recruitment period, 6146 patients were included (3580 women [58.2%]; median age, 71 [interquartile range (IQR), 63-76] years). Of these, 680 patients (11.1%) presented with postoperative complications. No differences were found in the number of patients with overall postoperative complications between ERAS and non-ERAS groups (163 [10.2%] vs 517 [11.4%]; odds ratio [OR], 0.89; 95% CI, 0.74-1.07; $P = .22$). Fewer patients in the ERAS group had moderate to severe complications (73 [4.6%] vs 279 [6.1%]; OR, 0.74; 95% CI, 0.56-0.96; $P = .02$). The median overall adherence rate with the ERAS protocol was 50.0% (IQR, 43.8%-62.5%), with the rate for ERAS facilities being 68.8% (IQR, 56.2%-81.2%) vs 50.0% (IQR, 37.5%-56.2%) at non-ERAS centers ($P < .001$). Among the patients with the highest and lowest quartiles of adherence to ERAS components, the patients with the highest adherence had fewer overall postoperative complications (144 [10.6%] vs 270 [13.0%]; OR, 0.80; 95% CI, 0.64-0.99; $P < .001$) and moderate to severe postoperative complications (59 [4.4%] vs 143 [6.9%]; OR, 0.62; 95% CI, 0.45-0.84; $P < .001$) and shorter median length of hospital stay (4 [IQR, 3-5] vs 5 [IQR, 4-6] days; OR, 0.97; 95% CI, 0.96-0.99; $P < .001$).

CONCLUSIONS AND RELEVANCE An increase in adherence to the ERAS program was associated with a decrease in postoperative complications, although only a few ERAS items were individually associated with improved outcomes.

JAMA Surg. doi:10.1001/jamasurg.2019.6024
Published online February 12, 2020.

Author Affiliations: Author affiliations are listed at the end of this article.

Group Information: Members of the POWER2 Study Investigators Group for the Spanish Perioperative Audit and Research Network appear at the end of the article.

Corresponding Author: Javier Ripollés-Melchor, MD, Department of Anaesthesia and Critical Care, Infanta Leonor University Hospital, Gran Vía del Este 80, 28031 Madrid, Spain (ripo542@gmail.com).

An estimated 310 million patients undergo surgery worldwide each year.¹ In the United States, 1 million total hip arthroplasty (THA) and total knee arthroplasty (TKA) procedures are performed annually, with an expectation of growth in the coming years.² The International Surgical Outcomes Study³ showed that 16.5% of patients who underwent orthopedic surgery experienced complications. The incidence of complications in THA and TKA was 3.2% to 8.0%,⁴ with an important variability between centers.^{4,5} Postoperative complications after THA and TKA have been shown to increase hospital length of stay (LOS).⁶ Kehlet⁷ described the concept of the multimodal approach to recovery in 1997. The Enhanced Recovery After Surgery (ERAS) program that entails an evidenced-based protocol based on a wide literature search was initiated by Ljungqvist et al⁸ in the ERAS study group, which later developed into the ERAS Society. The ERAS protocol includes a variable number of multidisciplinary perioperative measures whose initial goal is to improve recovery and reduce morbidity, which will result in secondary benefits, including reductions in LOS, convalescence, and costs.⁹ Observational studies¹⁰ and randomized clinical trials¹¹ have shown that ERAS was associated with improved outcomes in patients undergoing THA and TKA, mainly a decrease in LOS.^{12,13}

Until the recent publication of the ERAS Society THA and TKA guidelines,¹⁴ the most accepted recommendations were those proposed by Soffin and YaDeau in 2016.¹⁵ ERAS in THA and TKA share the same problems as ERAS in other surgical procedures: adoption is slow,¹⁶ and adherence to ERAS guidelines is low.¹⁷

The aim of this study was to characterize the management strategies for patients who underwent elective THA or TKA, including centers with and without an established ERAS protocol. The primary end point was the incidence of postoperative complications. Secondary end points were LOS and the potential associations between the adherence to the individual ERAS items and the occurrence of postoperative complications.

Methods

Study Design and Participants

This study is reported in accordance with the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline¹⁸ and the Reporting on ERAS Compliance, Outcomes, and Elements Research (RECOVER) Checklist.¹⁹ Hospital and investigator engagement was facilitated through the Spanish Perioperative Audit and Research Network (REDGERM), which was established by the Grupo Español de Rehabilitación Multimodal (GERM). All hospitals in Spain were invited to participate regardless of the number of beds and the existence of an ERAS protocol.

The Postoperative Outcomes Within Enhanced Recovery After Surgery Protocol in Elective Total Hip and Knee Arthroplasty (POWER2) study was a prospective 2-month multicenter cohort study. The study was approved by the Instituto Aragonés de Ciencias de la Salud Ethics Committee, Zaragoza, Spain, and the Spanish Medical Agency and was regis-

Key Points

Question How frequent are complications after total hip and knee arthroplasty procedures and how are the Enhanced Recovery After Surgery (ERAS) protocols associated with them?

Findings In this large, multicenter cohort study that included 6146 patients in 131 hospitals, 680 (11.1%) presented with postoperative complications and 352 (5.7%) presented with complications graded as moderate to severe. Patients who had greater compliance with the ERAS items had fewer postoperative complications, regardless of whether or not the center had an established ERAS protocol.

Meaning Although only a few perioperative interventions were associated with decreased complications, greater adherence with the set of ERAS measures was associated with a decrease in postoperative complications at 30 days of follow-up.

tered prospectively in an international trial registry.²⁰ The study protocol, technical appendix, and other documents are available at <http://www.grupogerm.es/power2>, and the study protocol has been published.²¹

The ethics committees or institutional review boards at each site reviewed and approved the protocol. Written informed consent was obtained from all participants.

Procedures

All consecutive adult patients (aged >18 years) undergoing elective primary THA or TKA were included during a single period of recruitment from October 22 through December 22, 2018, in each participating center. Each patient was followed up for 30 days after surgery. The follow-up was performed through hospital and primary care medical records.

Data were collected using Castor Electronic Data Capture²² and anonymized before entry onto a secure Internet-based electronic case record form designed specifically for POWER2. Participating hospitals were considered self-proclaimed ERAS centers in cases in which a multidisciplinary clinical pathway was implemented, regardless of the perioperative items that conformed to the pathway and whether hospitals were audited regularly. Individual data on 16 ERAS items were collected prospectively for each included patient. The definition of the individual ERAS components was based on the recommendations of Soffin and YaDeau¹⁵ (eTable 1 in the Supplement). Data included patient characteristics (American Society of Anesthesiologists [ASA] score, age, sex, smoking status, body mass index, clinical frailty score, and comorbidities), procedure performed, surgical approach, perioperative interventions, adherence to ERAS items, and outcomes (including postoperative complications, time to achieve targeted mobility, LOS, and 30-day mortality). Complications were defined and graded as mild, moderate, or severe according to international recommendations^{21,23-26} (eTable 2 in the Supplement) and were included if they occurred within 30 days after surgery. Data validation was conducted by specific validators at each site. A minimum sample size of 3012 was estimated, expecting 50% of patients with at least 1 complication, a 95% confidence level, and 3% accuracy.

Outcomes

The primary outcome was the number of patients with 30-day in-hospital postoperative complications. Secondary outcome measures included the number of patients with 30-day in-hospital postoperative moderate to severe complications, mortality, readmission, reintervention rates, and hospital LOS.

Statistical Analysis

Data were analyzed from March 15 through September 15, 2019. We analyzed outcomes depending on whether the patient received treatment at a hospital that belonged to a self-designated ERAS program (ERAS group). The discrete and continuous variables were described as number (percentage) and median (interquartile range [IQR]), respectively. We analyzed differences using the Fisher exact test for discrete variables and the Pearson and Wilcoxon rank sum tests for continuous variables. Subsequently, we repeated the analysis, subdividing the sample into quartiles according to the real adherence rate to ERAS items (regardless of whether the patient was treated at an ERAS hospital or not) and comparing the quartiles of higher and lower adherence to ERAS items. Next, we analyzed complication rates for each of the ERAS items using the Fisher exact test, and we performed a multivariable logistic regression analysis to study the association of the rate of each of the items with the clinical and demographic variables. We also used the same model in a multilevel multivariable logistic regression model to explore independent factors associated with postoperative complications assessing the variability of each site. Stratification by type of surgery (THA or TKA) was performed for the most relevant outcomes.


We performed several post hoc analyses. To assess the association of the underlying pathologic finding in the included patients, we divided the patients between those with ASA scores of I through II and III through IV (higher scores indicate severe preoperative systemic disease) and analyzed the adherence to ERAS items in these groups, as well as the postoperative outcomes. We performed this analysis for the self-described ERAS centers vs non-ERAS centers and for the ERAS adherence quartiles, depending on the median compliance in each group. Bubble plots were used to graphically illustrate the associations between the percentage of patients with postoperative complications and LOS with the volume of patients recruited at each site. To study the association between the percentage of patients with postoperative complications and LOS and the volume of patients recruited by each center, we used 2 β regression models. Through a linear regression model, we performed a post hoc analysis to evaluate the influence of ERAS items and other perioperative variables in LOS. For this purpose, we performed a univariable and a multivariable analysis. We denoted as statistically significant those comparisons where 2-sided $P < .05$.

Results

Participants

Data describing 6146 patients were collected in 131 hospitals in Spain (Figure 1). Of these, 2566 (41.8%) were men and

Figure 1. STROBE Flow Diagram for Included Patients



3580 were women (58.2%); median age was 71 (IQR, 63-76) years. Other characteristics are shown in Table 1. No centers or patients were excluded owing to low recruitment. According to the hospitals where the surgeries were performed, 1592 patients (25.9%) were included in self-designated ERAS centers. The ERAS and non-ERAS groups showed demographic differences in the number of patients with hypertension (2747 [60.3%] vs 900 [56.5%]), coronary artery disease (290 [6.4%] vs 70 [4.4%]), atrial fibrillation (336 [7.4%] vs 85 [5.3%]), chronic kidney disease (224 [4.9%] vs 108 [6.8%]), ASA score (ASA III-IV (345 [21.7%] vs 1273 [28.0%]), and duration of surgery (median, 90 [IQR, 75-115] vs 83 [IQR, 65-104] minutes). More patients in the ERAS group received local anesthesia (188 [11.8%] vs 64 [1.4%]) and tranexamic acid (1348 [85.9%] vs 3191 [70.2%]), whereas fewer patients in the ERAS group received epidural (38 [2.4%] vs 350 [7.7%]) or regional (157 [9.9%] vs 770 [16.9%]) anesthesia. Patients in the ERAS group also received fewer crystalloids (median, 800 [IQR, 550-1000] vs 1000 [IQR, 700-1200] mL) and colloids (median, 0 [IQR, 0-250] vs 0 [IQR, 0-0] mL) during surgery as well as fewer red blood cell transfusions (RBCTs) (mean [SD], 9 [0.6%] vs 52 [1.1%]) (Table 1). ERAS adherence quartiles were calculated for all patients from the median overall adherence to the 16 ERAS items, regardless of whether the patient was treated within an ERAS protocol. Adherence in quartile 1 was 43.75% or less; in quartile 2, greater than 43.75% to 50.00%; in quartile 3, greater than 50.00% to 62.50%; and in quartile 4, greater than 62.50%. The quartile with the lowest adherence to ERAS included the most patients (eTable 3 in the Supplement). Patients with lower adherence to ERAS had a higher prevalence of chronic kidney disease (48 [4.4%] vs 103 [5.0%]), had longer duration of surgery (median, 90 [IQR, 74-110] vs 90 [IQR, 75-115] minutes), received more crystalloid during surgery (median, 1000 [IQR, 700-1200] vs 1000 [750-1200] mL), and had more epidural anesthesia (73 [6.8%] vs 149 [7.2%]) and regional analgesia (185 [17.1%] vs 244 [11.8%]) performed compared with the group with greater adherence to ERAS (eTable 3 in the Supplement).

Table 1. Demographic and ERAS Adherence Data^a

Characteristic	Patient Group		
	All (n = 6146)	No ERAS (n = 4554)	ERAS (n = 1592)
Age, median (IQR), y	71 (63-76)	71 (64-76)	70 (63-76)
Men	2566 (41.8)	1904 (41.8)	662 (41.6)
BMI, median (IQR)	29.3 (26.4-32.9)	29.4 (26.4-32.8)	29.3 (26.4-32.9)
ASA classification ^b			
ASA I	391 (6.4)	262 (5.8)	129 (8.1)
ASA II	4136 (67.3)	3019 (66.3)	1117 (70.2)
ASA III	1575 (25.6)	1238 (27.2)	337 (21.2)
ASA I-II	4257 (73.7)	3281 (72.0)	1246 (78.3)
ASA III-IV	1618 (26.3)	1273 (28.0)	345 (21.7)
Former smoker	1005 (16.4)	758 (16.6)	247 (15.5)
Current smoker	699 (11.4)	505 (11.1)	194 (12.2)
Frailty score, median (IQR) ^c	3 (2-3)	3 (2-3)	3 (2-4)
Hypertension	3647 (59.3)	2747 (60.3)	900 (56.5)
Diabetes	1140 (18.5)	848 (18.6)	292 (18.3)
Coronary arterial disease	360 (5.9)	290 (6.4)	70 (4.4)
Heart failure	158 (2.6)	113 (2.5)	45 (2.8)
Cirrhosis	52 (0.8)	42 (0.9)	10 (0.6)
Stroke	288 (4.7)	226 (5.0)	62 (3.9)
Atrial fibrillation	421 (6.8)	336 (7.4)	85 (5.3)
COPD/asthma	694 (11.3)	522 (11.5)	172 (10.8)
Chronic kidney disease	332 (5.4)	224 (4.9)	108 (6.8)
Peripheral vasculopathy	281 (4.6)	217 (4.8)	64 (4.0)
Rheumatoid arthritis	126 (2.1)	94 (2.1)	32 (2.0)
Dementia	66 (1.1)	55 (1.2)	11 (0.7)
Hemoglobin level, median (IQR), g/dL	14.0 (13.1-15.0)	14.1 (13.1-15.0)	14.0 (13.1-15.0)
Creatinine level, median (IQR), mg/dL	0.80 (0.70-0.95)	0.80 (0.69-0.95)	0.80 (0.70-0.96)
THA	2301 (37.4)	1711 (37.6)	590 (37.1)
TKA	3845 (62.6)	2843 (62.4)	1002 (62.9)
Duration of surgery, median (IQR), min	90 (75-110)	90 (75-115)	83 (65-104)
Anesthesia			
General	460 (7.5)	351 (7.7)	109 (6.8)
Intradural	5679 (92.4)	4202 (92.3)	1477 (92.8)
Epidural	388 (6.3)	350 (7.7)	38 (2.4)
Regional	927 (15.1)	770 (16.9)	157 (9.9)
Local	252 (4.1)	64 (1.4)	188 (11.8)
Crystalloid administered, median (IQR), mL	1000 (700-1200)	1000 (700-1200)	800 (550-1000)
Colloid administered, median (IQR), mL	0 (0-0)	0 (0-0)	0 (0-250)
Red blood cell transfusion	61 (1.0)	52 (1.1)	9 (0.6)
Red blood cell transfused, median (IQR), mL	445 (260-560)	400 (258-500)	715 (525-800)
Blood loss, median (IQR), mL	200 (100-350)	200 (100-350)	200 (100-400)
Tranexamic acid administered	4539 (73.9)	3191 (70.2)	1348 (85.9)
Presurgical education	1688 (27.5)	433 (9.5)	1255 (78.8)
Presurgical optimization	1193 (19.4)	547 (12.0)	646 (40.6)
Preoperative fasting	2997 (48.8)	1831 (40.2)	1166 (73.2)
Patient blood management	2396 (39.0)	1273 (28.0)	1123 (70.5)
Preoperative carbohydrate drinks preload	117 (1.9)	78 (1.7)	39 (2.4)
Avoidance of long-acting sedative premedication	3311 (53.9)	2448 (53.8)	863 (54.2)
Thromboprophylaxis	5655 (92.0)	4120 (90.5)	1535 (96.4)
Antibiotic prophylaxis	6076 (98.9)	4497 (98.7)	1579 (99.2)
Regional anesthesia	4919 (80.0)	3575 (78.5)	1344 (84.4)

(continued)

Table 1. Demographic and ERAS Adherence Data^a (continued)

Characteristic	Patient Group		
	All (n = 6146)	No ERAS (n = 4554)	ERAS (n = 1592)
PONV prophylaxis	3842 (62.5)	2739 (60.1)	1103 (69.3)
Active prevention of unintentional hypothermia	5082 (82.7)	3734 (82.0)	1348 (84.7)
Goal-directed fluid therapy	591 (9.6)	269 (5.9)	322 (20.2)
Postoperative analgesia	5978 (97.3)	4410 (96.8)	1568 (98.5)
Postoperative glycemic control	3208 (52.2)	2258 (49.6)	950 (59.7)
Early mobilization	2166 (35.2)	984 (21.6)	1182 (74.2)
Early feeding	3953 (64.3)	2563 (56.3)	1390 (87.3)

Abbreviations: ASA, American Society of Anesthesiologists; BMI, body mass index (calculated as weight in kilograms divided by height in meters squared); COPD, chronic obstructive pulmonary disease; ERAS, Enhanced Recovery After Surgery; IQR, interquartile range; PONV, postoperative nausea and vomiting; THA, total hip arthroplasty; TKA, total knee arthroplasty.

SI conversion factors: To convert albumin to grams per liter, multiply by 10; creatinine to micromoles per liter, multiply by 88.4; hemoglobin to grams per liter, multiply by 10.

^a A complete definition list of all ERAS items was published in the protocol.¹⁷ Unless otherwise indicated, data are expressed as number (percentage) of patients.

^b Higher scores indicate severe preoperative systemic disease.

^c Scores range from 1 to 9, with higher scores indicating higher level of vulnerability.

Outcome Data

A total of 680 patients (11.1%) experienced overall postoperative complications, and 352 (5.7%) experienced complications graded as moderate to severe (eFigure 1 in the Supplement). No differences were found in the number of patients with overall postoperative complications between the ERAS and non-ERAS groups (163 [10.2%] vs 517 [11.4%]; OR, 0.89; 95% CI, 0.74-1.07; $P = .22$) or in terms of readmission, reoperation, or survival (eFigure 1 in the Supplement). Fewer patients in the ERAS group had moderate to severe complications (73 [4.6%] vs 279 [6.1%]; OR, 0.74; 95% CI, 0.56-0.96; $P = .02$). eTables 4 and 5 in the Supplement show the stratified complications in THA and TKA in ERAS vs non-ERAS centers. In THA and TKA, the ERAS group had shorter median LOS (THA, 4 [IQR, 3-5] vs 5 [IQR, 4-6] days; TKA, 4 [IQR, 3-5] vs 5 [IQR, 4-6] days) (eFigure 1 and eTables 4 and 5 in the Supplement). Graduation of the severity of postoperative complications is shown in eTable 6 of the Supplement.

ERAS Adherence and Outcome Data

The overall adherence rate to the ERAS protocol components was 50.0% (IQR, 43.8%-62.5%), with the rate for ERAS facilities being 68.8% (IQR, 56.2%-81.2%) vs 50.0% (IQR, 37.5%-56.2%) at non-ERAS centers ($P < .001$). Adherence to specific ERAS items was greater in the ERAS group (Table 1). The ERAS group also had less bleeding (median blood loss, 200 [IQR, 100-400] vs 200 [IQR, 100-350] mL), less use of intraoperative crystalloids (median, 800 [IQR, 550-1000] vs 1000 [IQR, 700-1200] mL), and greater intraoperative administration of tranexamic acid (1348 [85.9%] vs 3191 [70.2%]) (Table 1). In the TKA and THA procedures, adherence to perioperative ERAS items was greater in the ERAS group (eTables 7, 8, and 9 in the Supplement).


Patients in the ERAS group undergoing THA and TKA had less time to ambulation (mean [SD] for THA, 17.2 [13.3] vs 30.9 [24.4] hours; mean [SD] for TKA, 15.2 [13.9] vs 29.1 [15.5] hours) and oral intake (mean [SD] for THA, 6.04 [5.12] vs 8.12 [11.2] hours; mean [SD] for TKA, 6.05 [5.11] vs 8.20 [6.07] hours)

(eTable 10 in the Supplement). Among the highest (quartile 4) and lowest (quartile 1) quartiles of adherence to ERAS items, quartile 4 also had a decrease in the rate of overall postoperative complications (270 [13.0%] vs 144 [10.6%]; OR, 0.80; 95% CI, 0.64-0.99; $P < .001$), moderate to severe postoperative complications (143 [6.9%] vs 59 [4.4%]; OR, 0.62; 95% CI, 0.45-0.84; $P < .001$), postoperative hemorrhage (106 [5.1%] vs 31 [2.3%]; OR, 0.44; 95% CI, 0.29-0.65; $P < .001$), infection of uncertain source (17 [0.8%] vs 3 [0.2%]; OR, 0.28; 95% CI, 0.06-0.84; $P = .02$), and surgical complications (37 [1.8%] vs 12 [0.9%]; OR, 0.50; 95% CI, 0.25-0.93; $P = .03$) (Figure 2). Graduation of the severity of postoperative complications is shown in eTable 11 in the Supplement. eFigures 2 and 3 in the Supplement show the results grouped by THA and TKA. In THA and TKA, the highest ERAS adherence group had shorter LOS (eFigures 2 and 3 in the Supplement) and less time to ambulation and oral intake (eFigure 4 in the Supplement) in THA and TKA (eFigures 5 and 6 in the Supplement).

Multivariable analysis of the associations with ERAS items showed a statistically significant reduction of complications in patients who had regional anesthesia (OR, 0.71; 95% CI, 0.53-0.96; $P = .02$) and early mobilization (OR, 0.61; 95% CI, 0.45-0.83; $P = .002$) (Table 2). A higher preoperative hemoglobin value (OR, 0.84; 95% CI, 0.77-0.91; $P < .001$) and tranexamic acid administration (OR, 0.66; 95% CI, 0.51-0.86; $P = .002$) were associated with fewer complications, whereas blood loss was associated with postoperative complications (OR, 1.00; 95% CI, 1.00-1.00; $P < .001$) (Table 2). In the multilevel study, early mobilization was associated with decreased complications (OR, 0.60; 95% CI, 0.42-0.86; $P = .005$) but not regional anesthesia (OR, 0.75; 95% CI, 0.54-1.05; $P = .10$) (Table 2).

Stratification of adherence to individual ERAS items by severity groups showed that healthier patients had higher adherence to ERAS (eTable 12 in the Supplement). In the ERAS centers, patients with ASA scores of I to II experienced less infection of uncertain source (1 [0.1%] vs 16 [0.5%]; OR, 0.19; 95% CI, 0.01-0.91; $P = .04$), less hemorrhage (18 [0.1%] vs 104 [3.2%]; OR, 0.45; 95% CI, 0.26-0.73; $P = .001$), and lower median LOS

Figure 2. Postoperative Outcomes and Enhanced Recovery After Surgery (ERAS) Adherence



Patients are stratified by complications, type of complication, and other secondary outcomes and by quartile of adherence to the ERAS protocol. Surgical complications included all postoperative complications as defined by protocol¹⁷ according to the standards for definitions and use of outcome measures for clinical effectiveness research in perioperative medicine: European Perioperative Clinical Outcome (EPCO) definitions,²³ the standardized list, definitions, and the stratification developed by the Hip

Society,²⁴ the standardized list and definitions of the Knee Society,²⁵ and the Standardized Bleeding Definitions for Cardiovascular Clinical Trials.²⁶ IQR indicates interquartile range; LOS; length of stay; and OR, odds ratio.

^a Unless otherwise indicated, data are expressed as number (percentage) of patients.

(4 [IQR, 3-5] vs 5 [IQR, 3-6] days; OR, 0.95; 95% CI, 0.94-0.97; $P < .001$) compared with the non-ERAS centers (eFigure 7 in the Supplement), whereas patients with ASA scores of III to IV in the ERAS centers experienced less readmission compared with non-ERAS centers (18 [5.2%] vs 36 [28%]; OR, 1.90; 95% CI, 1.04-3.35; $P = .04$) (eFigure 8 in the Supplement). Regardless of belonging to a ERAS center or not, patients with ASA scores of I to II with greater adherence to ERAS had fewer moderate to severe complications (31 [3.0%] vs 79 [5.3%]; OR, 0.55; 95% CI, 0.35-0.83; $P = .004$) and lower median LOS (4 [IQR, 3-5] vs 5 [IQR, 4-6] days; OR, 0.97; 95% CI, 0.95-0.98; $P < .001$) compared with those with less adherence to ERAS (eFigure 9 in the Supplement), whereas no difference was found between patients with ASA scores of III to IV with more or less adherence to ERAS (eFigure 10 in the Supplement).

No significant association was found among the volume of patients recruited, hospital LOS, and postoperative complications. LOS decreased a mean of 0.0048 days due to an increase in 1 patient ($P = .20$, associated with β regression models), whereas the increase of 1 person per center increased the postoperative complications by 0.37% ($P = .10$, associ-

ated with β regression models) (Figure 3 and eFigures 11-15 in the Supplement).

Median LOS was shorter in the ERAS group compared with the non-ERAS group (4 [IQR, 3-5] vs 5 [IQR, 4-6] days; $P < .001$). Regarding the difference by adherence quartiles, quartile 1 presented with a median LOS of 5 (IQR, 4-6) days, compared with 4 (IQR, 3-5) days in the highest adherence group (OR, 0.97; 95% CI, 0.96-0.99; $P < .001$) (Figure 3). The post hoc multivariable analysis of the influence of patient characteristics and ERAS items showed a statistically significant reduction of LOS in patients who achieved early mobilization (β coefficient, -0.51 ; 95% CI, -0.95 to -0.06 ; $P = .03$) and received local anesthesia (β coefficient, -1.59 ; 95% CI, -2.61 to -0.57 ; $P = .002$), whereas active prevention of unintentional hypothermia (β coefficient, 0.78 ; 95% CI, 0.30 - 1.26 ; $P = .001$), epidural anesthesia (β coefficient, 1.53 ; 95% CI, 0.79 - 2.26 ; $P < .001$), and complications (β coefficient, 2.63 ; 95% CI, 2.00 - 3.25 ; $P < .001$) were associated with an increase in LOS (eTable 13 in the Supplement). It was not necessary to censor any P value after the multiple comparison study to adjust the false discovery rate to 5%.

Table 2. Multivariable and Multilevel Analysis for Postoperative Complications^a

Variable	No Complications	Complications	Univariable Analysis		Multivariable Analysis		Multilevel Analysis	
			OR (95% CI)	P Value	OR (95% CI)	P Value	OR (95% CI)	P Value
Age, mean (SD), y	68.8 (10.2)	70.8 (10.5)	1.02 (1.01-1.03)	<.001	1.00 (0.99-1.02)	.46	1.00 (0.99-1.02)	.64
BMI, mean (SD)	30.1 (6.3)	30.0 (6.4)	1.00 (0.98-1.01)	.68	0.99 (0.97-1.01)	.35	0.99 (0.97-1.01)	.31
Frailty score, mean (SD) ^b	3.0 (1.0)	3.3 (1.1)	1.34 (1.25-1.44)	<.001	1.02 (0.89-1.16)	.75	1.04 (0.90-1.20)	.58
ASA classification ^c								
ASA I	360 (92.1)	31 (7.9)	NA	NA	NA	NA	NA	NA
ASA II	3765 (91.0)	371 (9.0)	1.14 (0.79-1.71)	.49	1.22 (0.66-2.47)	.56	1.26 (0.64-2.48)	.50
ASA III	1309 (83.1)	266 (16.9)	2.36 (1.62-3.55)	<.001	2.14 (1.11-4.50)	.03	2.23 (1.08-4.59)	.03
ASA IV	31 (72.1)	12 (27.9)	4.50 (2.04-9.47)	<.001	2.41 (0.68-8.02)	.16	3.07 (0.84-11.22)	.09
Diabetes	165 (82.1)	36 (17.9)	1.80 (1.22-2.57)	.002	1.10 (0.59-1.96)	.74	1.03 (0.55-1.95)	.924
Heart failure	122 (77.2)	36 (22.8)	2.45 (1.65-3.54)	<.001	1.35 (0.72-2.42)	.33	1.40 (0.74-2.63)	.30
Atrial fibrillation	344 (81.7)	77 (18.3)	1.90 (1.46-2.46)	<.001	1.30 (0.86-1.93)	.20	1.31 (0.86-2.00)	.22
Chronic kidney disease	273 (82.2)	59 (17.8)	1.81 (1.34-2.41)	<.001	1.19 (0.72-1.91)	.48	1.19 (0.71-1.99)	.50
Hemoglobin level, mean (SD), g/dL	14.1 (1.4)	13.6 (1.5)	0.79 (0.75-0.84)	<.001	0.84 (0.77-0.91)	<.001	0.83 (0.76-0.91)	<.001
Blood loss, mean (SD), mL	248.6 (213.7)	301.5 (263.6)	1.00 (1.00-1.00)	<.001	1.00 (1.00-1.00)	<.001	1.00 (1.00-1.00)	<.001
Tranexamic acid administration	4111 (90.6)	428 (9.4)	0.56 (0.47-0.66)	<.001	0.66 (0.51-0.86)	.002	0.75 (0.56-1.00)	.048
Presurgical education	1509 (89.4)	179 (10.6)	0.94 (0.78-1.12)	.49	1.38 (0.96-1.98)	.08	1.31 (0.85-2.02)	.22
Presurgical optimization	1078 (90.4)	115 (9.6)	0.94 (0.75-1.18)	.62	0.86 (0.62-1.20)	.39	0.94 (0.64-1.39)	.76
Preoperative fasting	2683 (89.5)	314 (10.5)	0.89 (0.76-1.04)	.15	1.07 (0.82-1.40)	.61	1.06 (0.76-1.47)	.73
Patient blood management	2144 (89.5)	252 (10.5)	0.92 (0.78-1.08)	.29	0.93 (0.71-1.23)	.63	0.85 (0.61-1.19)	.35
Preoperative carbohydrate drinks preload	110 (94.0)	7 (6.0)	0.51 (0.21-1.01)	.08	0.72 (0.24-1.70)	.49	0.44 (0.14-1.37)	.16
Avoidance of long-acting sedative premedication	2945 (88.9)	366 (11.1)	1.00 (0.85-1.17)	.97	1.03 (0.80-1.32)	.82	1.11 (0.83-1.47)	.49
Thromboprophylaxis	5023 (88.8)	632 (11.2)	1.15 (0.85-1.58)	.39	1.04 (0.68-1.67)	.86	1.14 (0.70-1.87)	.60
Regional anesthesia	4406 (89.6)	513 (10.4)	0.74 (0.61-0.89)	.001	0.71 (0.53-0.96)	.02	0.75 (0.54-1.05)	.10
PONV prophylaxis	3427 (89.2)	415 (10.8)	0.93 (0.79-1.10)	.39	0.93 (0.72-1.21)	.59	0.90 (0.67-1.21)	.47
Active prevention of unintentional hypothermia	4506 (88.7)	576 (11.3)	1.17 (0.94-1.47)	.16	1.54 (1.11-2.18)	.01	1.61 (1.05-2.46)	.03
Goal-directed fluid therapy	539 (91.2)	52 (8.8)	0.76 (0.56-1.01)	.07	0.98 (0.64-1.47)	.97	0.87 (0.54-1.40)	.57
Postoperative analgesia	5313 (88.9)	665 (11.1)	1.28 (0.77-2.28)	.37	2.40 (1.04-6.99)	.07	2.28 (0.84-6.16)	.11
Postoperative glycemic control	2849 (88.8)	359 (11.2)	1.03 (0.88-1.20)	.74	0.79 (0.61-1.02)	.07	0.91 (0.68-1.22)	.53
Early mobilization	1985 (91.6)	181 (8.4)	0.64 (0.53-0.76)	<.001	0.61 (0.45-0.83)	.002	0.60 (0.42-0.86)	.005
Early feeding	3536 (89.5)	417 (10.5)	0.86 (0.73-1.02)	.08	1.13 (0.86-1.48)	.38	1.15 (0.85-1.56)	.36
Analgesia								
Epidural anesthesia	342 (88.1)	46 (11.9)	1.09 (0.78-1.48)	.61	1.06 (0.67-1.64)	.79	0.95 (0.57-1.57)	.83
Regional anesthesia	836 (90.2)	91 (9.8)	0.86 (0.67-1.07)	.19	1.03 (0.71-1.45)	.88	0.91 (0.60-1.38)	.67
Local anesthesia	230 (91.3)	22 (8.7)	0.76 (0.47-1.16)	.23	0.85 (0.36-1.72)	.76	1.05 (0.45-2.47)	.91

Abbreviations: ASA, American Society of Anesthesiologists; BMI, body mass index (calculated as weight in kilograms divided by square of height in meters); IQR, interquartile range; NA, not applicable; OR, odds ratio; PONV, postoperative nausea and vomiting.


SI conversion factors: To convert albumin to grams per liter, multiply by 10; creatinine to micromoles per liter, multiply by 88.4; hemoglobin to grams per liter, multiply by 10.

^a Statistically significant. Unless otherwise indicated, data are expressed as number (percentage) of patients.

^b Scores range from 1 to 9, with higher scores indicating higher level of vulnerability.

^c Higher scores indicate severe preoperative systemic disease.

Figure 3. Bubble Plot of the Rate of Postoperative Complications vs Length of Stay, Number of Participating Patients, and Rate of ERAS Program Adherence



For each hospital, the plot shows the mean percentage of adherence to the Enhanced Recovery After Surgery (ERAS) program (bubble color), the mean length of hospital stay, and percentage of patients who have postoperative complications. The bubble diameter indicates the number of patients that the hospital contributed to the study.

Discussion

POWER2 examined the postoperative outcomes in patients scheduled for THA or TKA and the outcomes associated with 16 evidence-based care components of an ERAS protocol in 131 centers in Spain and more than 6000 patients without limiting the inclusion of patients to centers with an established ERAS protocol. Results demonstrated that the implementation of an ERAS protocol was associated with improved postoperative outcomes.

A 2018 meta-analysis²⁷ found that ERAS protocols were associated with a significant decrease in transfusions, complication rates, and LOS without decreasing mortality or increasing the 30-day readmission rate in patients undergoing THA or TKA. During the last decade, the clinical pathways and care programs in THA and TKA have undergone considerable changes in many countries influenced by the ERAS concept.²⁸ However, in the absence of guidelines, the items that com-

prise these ERAS protocols were not clearly defined and present a great variability in the studies performed.²⁷ POWER2 found that the adherence to ERAS protocol in THA and TKA was low in all patients, although it was higher and close to 70% in self-proclaimed ERAS centers. The low adherence to ERAS protocols may be owing to the usual barriers to the implementation of ERAS in other disciplines, such as resistance to change, inadequate financing, lack of administration support, high turnover, poor documentation, and lack of time,²⁹ to which orthopedic procedures add the lack of guidelines and the economic pressure to achieve decreases in LOS.

An adherence of greater than 62.50% was associated with a decrease in postoperative complications, including mild and moderate to severe. Importantly, only some of the perioperative interventions seemed to be associated with a decrease in complications. Similarly, Khan et al¹² demonstrated that the application of an ERAS protocol based on a few components that included a multidisciplinary strategy involving patient educa-

tion, multimodal analgesia, standardized perioperative anesthesia, local anesthetic infiltration, judicious fluid administration, tranexamic acid administration, and early mobilization led to a reduction in complications, LOS, and 30-day mortality. Khan et al¹² showed mean hospital stays of 3 days for patients in ERAS centers, whereas in POWER2, the best results were a median of 4 days in the patients with greater adherence to ERAS. Although some centers and health systems have stays of 3 or fewer days and even perform THA and TKA as outpatient procedures, the usual LOS³⁰ is similar to that reported in POWER2. This may be because, in Spain, patients are directly discharged home. In contrast, postacute care is typically available and covered by insurance companies in the United States and in other countries, making it easy for hospitals to discharge patients to rehabilitation facilities or to other intermediate centers.³¹

POWER2 found no association between comorbidities or frailty and complications, although patients with an ASA physical status score of III had more complications. We found that in the ERAS centers and in the group with greater ERAS adherence, there were more healthy patients. This finding could explain the best results in these groups. In fact, although the ERAS programs in THA and TKA have been proven safe and beneficial for most patients, we have found that in high-risk patients (ASA scores of III-IV), the application of ERAS, even in the highest adherence quartiles, did not result in better outcomes, although there were fewer readmissions.¹²

In patients undergoing major orthopedic surgical procedures, preoperative anemia, perioperative bleeding, and a liberal transfusion policy are the main risk factors for requiring an RBCT,³² which increases complications and LOS.^{33,34} Yang et al³⁵ showed that in patients undergoing TKA, tranexamic acid administration reduced blood loss and the need for RBCT. Similarly, we found that lower preoperative hemoglobin values and intraoperative bleeding were associated with an increase in complications, whereas the administration of tranexamic acid was associated with a decrease. Avoiding anemia, improving tolerance to anemia, and avoiding bleeding and RBCT are the pillars of patient blood management programs.³⁶ Surprisingly, the ERAS patient blood management item included in POWER2 has not been associated with better outcomes. In our opinion, many centers may apply the patient blood management measures without recognizing them as such.

It is not surprising that good analgesic control leads to early mobilization, and accomplishment of these items was associated with improved outcomes. Early mobilization in the first 24 hours after THA or TKA had been shown to be effective in reducing LOS, although it had not been associated with a decrease in complications.³⁷ Nevertheless, in POWER2, early mobilization was associated with fewer complications and shorter LOS. Conversely, epidural anesthesia was associated with a greater than 1-day increase in hospital LOS.

Early mobilization is considered a cornerstone in ERAS pathways and is strongly recommended as part of ERAS guidelines in many surgical disciplines.⁸ ERAS Society guidelines recommend that patients be mobilized as soon as possible after surgery.¹⁴ Chua et al³⁸ found that for patients undergoing THA, shorter time to mobilization was associated with hospital site and the absence of acute complications. For patients under-

going TKA, shorter time to mobilization was associated with hospital site and absence of donor blood transfusion.³⁸ These outcomes indicate that the patient who has complications cannot move early, and delays in mobilization could be taken as an alarm signal for possible complications. However, the delay until mobilization can also be related to the institution and local protocols, because certain hospitals encourage early patient mobilization, whereas others do not.

Strengths and Limitations

POWER2 had several strengths compared with other studies. This was a prospective study, with a case report form designed specifically for the study and with a short recruitment period of 2 months. We also included a large number of centers. Other studies have had years-long durations.³⁴ Khan et al¹² showed that the application of an ERAS program improved outcomes in a single center and in a long period of 7 years; the changes produced in such long periods decrease patient comparability. On the other hand, some of the studies with the highest number of cases were registers in national databases, in which it is assumed that all patients were treated with an ERAS protocol, but in which the adherence was not analyzed. We have included self-described ERAS centers and non-ERAS centers, and we have analyzed the variables that make up the ERAS protocol in all the centers, unlike the studies based on databases or the before-and-after studies. Moreover, the inclusion of consecutive, nonselected patients from 131 different centers without any selection bias increases the internal validity of our study. In addition, there were few losses of follow-up.

Our study has certain limitations. It was observational, which means that we can only obtain associations. Second, as previously mentioned, early mobilization can act as a confounding factor; because patients with complications could not fulfill this item, whether mobilization is an exposure factor or an outcome is debatable. We believe that it must be treated as both, and that all patients should be encouraged to move as soon as possible after surgery, because this has been shown to improve postoperative outcomes.³⁹ Third, we chose the individual ERAS items based on expert recommendations¹⁵; if we had selected other items, the results might be different. On the other hand, the patients in the ERAS group probably corresponded to more specialized centers and were healthier compared with patients in non-ERAS centers, which could explain why they had better outcomes.

Conclusions

This study found that ERAS protocols in THA and TKA do not appear to be applied in daily clinical practice, even in self-proclaimed ERAS centers, even though a greater adherence to ERAS improves postoperative outcomes. An effective pathway for THA and TKA consists of regional or local analgesia, anemia and bleeding management, and early mobilization as the basis of the care trajectory. Consequently, our findings suggest that the focus should be on these essential elements of care rather than on expanding adherence to the other, less evidence-based components in most ERAS protocols.

ARTICLE INFORMATION

Accepted for Publication: December 1, 2019.

Published Online: February 12, 2020.

doi:10.1001/jamasurg.2019.6024

Author Affiliations: Department of Anaesthesia and Critical Care, Infanta Leonor University Hospital, Madrid, Spain (Ripollés-Melchor, Abad-Motos, Abad-Gurumeta); School of Medicine, Universidad Complutense de Madrid, Madrid, Spain (Ripollés-Melchor, Abad-Motos, Abad-Gurumeta); Spanish Perioperative Audit and Research Network (REDGERM) (Ripollés-Melchor, Abad-Motos, Suárez-de-la-Rica, Ferrando-Ortolá, Ramírez-Rodríguez, García-Erce, Aldecoa); Grupo Español de Rehabilitación Multimodal, Enhanced Recovery After Surgery Spain Chapter, Madrid (Ripollés-Melchor, Abad-Motos, Suárez-de-la-Rica, Cuéllar-Martínez, Ramiro-Ruiz, Abad-Gurumeta, Ferrando-Ortolá, Ramírez-Rodríguez, García-Erce, Aldecoa); Department of Anaesthesia and Critical Care, Ramón y Cajal University Hospital, Madrid, Spain (Diez-Remesal); Department of Anaesthesia, Donostia University Hospital/Donostia Unibertsitate Ospitalea, San Sebastián, Spain (Aseguinolaza-Pagola); Department of Anaesthesia, Complejo Hospitalario Universitario De Vigo-Eoxi Vigo, Vigo, Spain (Padin-Barreiro); Department of Anaesthesia, Hospital Universitario Clínico San Carlos, Madrid, Spain (Sánchez-Martín); Department of Anaesthesia, Hospital Universitario de Álava, Vitoria-Gasteiz, Spain (Logroño-Egea); Department of Anaesthesia, Consorcio Hospital General Universitario de Valencia, Valencia, Spain (Catalá-Bauset); Department of Anaesthesia, Hospital Universitario Sierrallana, Torrelavega, Spain (García-Orallo); Department of Anaesthesia, Hospitales del Parc de Salut Mar, Barcelona, Spain (Bisbe); Department of Anaesthesia, Hospital Clínic de Barcelona, Barcelona, Spain (Martín, Ferrando-Ortolá); Department of Anaesthesia and Critical Care, La Paz University Hospital, Madrid, Spain (Suárez-de-la-Rica); Department of Anaesthesia and Critical Care, Hospital Universitario Central de Asturias, Oviedo, Spain (Cuéllar-Martínez); Department of Anaesthesia, Hospital General Universitario de Ciudad Real, Ciudad Real, Spain (Gil-Trujillo); Department of Anaesthesia, Hospital Universitario Rey Juan Carlos, Móstoles, Spain (Estupiñán-Jiménez); Department of Anaesthesia, Hospital de Mataró, Mataró, Spain (Villanova-Baraza); Department of Anaesthesia, Hospital Universitario de Fuenlabrada, Fuenlabrada, Spain (Gil-Lapetra); Department of Anaesthesia, Hospital de Galdakao-Usansolo, Bizkaia, Spain (Pérez-Sánchez); Department of Anaesthesia, Hospital de La Cruz Roja, Gijón, Spain (Rodríguez-García); Department of Anaesthesia, Hospital Universitario 12 de Octubre, Madrid, Spain (Ramiro-Ruiz); Department of Anaesthesia, Althaia Xarxa Assistencial Universitèria de Manresa, Barcelona, Spain (Farré-Tebar); Department of Anaesthesia, Hospital Universitario de Jaén, Jaén, Spain (Martínez-García); Department of Anaesthesia, Hospital Quirón, Zaragoza, Spain (Arauzo-Pérez); Department of Anaesthesia, Hospital Universitario de León, León, Spain (García-Pérez); Department of Anaesthesia, Hospital Clínico Universitario de Valladolid, Valladolid, Spain (Miñambres-Villar); Department of Anaesthesia, Hospital Universitario de Cruces, Bilbao, Spain (Sánchez-Campos); Department of

Anaesthesia, Hospital Universitario Virgen del Rocío, Sevilla, Spain (Jiménez-López); Department of Anaesthesia, Complejo Hospitalario de Mérida, Mérida, Spain (Tena-Guerrero); Department of Orthopedía, Infanta Leonor University Hospital, Madrid, Spain (Marín-Peña); Department of Anaesthesia, Hospital Universitario Fundación Alcorcón, Alcorcón, Spain (Sánchez-Merchant); Department of Anaesthesia, Clínica Santa Cristina, Albacete, Spain (Vicente-Gutiérrez); Department of Anaesthesia, Hospital Universitario Miguel Servet, Zaragoza, Spain (Cassinello-Ogea); Department of Anaesthesia, Hospital Universitario Virgen Macarena, Sevilla, Spain (Berges-Gutiérrez); Department of Anaesthesia, Hospital Sant Joan Despí Moisès Broggi, Sant Joan Despí, Spain (Fernanz-Antón); Department of Anaesthesia, Complejo Hospitalario Universitario de A Coruña, Coruña, Spain (Gómez-Ríos); Instituto Aragonés de Ciencias de la Salud, Zaragoza, Spain (Bordonaba-Bosque, Ramírez-Rodríguez); Department of Surgery, Lozano Blesa University Hospital, Zaragoza, Spain (Ramírez-Rodríguez); Blood and Tissue Bank of Navarra, Servicio Navarro de Salud-Osasunbidea, Pamplona, Navarra, Spain (García-Erce); Department of Anaesthesia and Perioperative Medicine, Río Hortega University Hospital, Valladolid, Spain (Aldecoa).

Author Contributions: Dr Ripollés-Melchor had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Ripollés-Melchor, Abad-Motos, García-Orallo, Ferrando-Ortolá, Ramírez-Rodríguez, García-Erce, Aldecoa.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Ripollés-Melchor, Padin-Barreiro, Rodríguez-García, Ferrando-Ortolá, Bordonaba-Bosque.

Critical revision of the manuscript for important intellectual content: Ripollés-Melchor, Abad-Motos, Díez-Remesal, Aseguinolaza-Pagola, Sánchez-Martín, Logroño-Egea, Catalá-Bauset, García-Orallo, Bisbe, Martín, Suárez-de-la-Rica, Cuéllar-Martínez, Gil-Trujillo, Estupiñán-Jiménez, Villanova-Baraza, Gil-Lapetra, Pérez-Sánchez, Ramiro-Ruiz, Farré-Tebar, Martínez-García, Arauzo-Pérez, García-Pérez, Abad-Gurumeta, Miñambres-Villar, Sánchez-Campos, Jiménez-López, Tena-Guerrero, Marín-Peña, Sánchez-Merchant, Vicente-Gutiérrez, Cassinello-Ogea, Ferrando-Ortolá, Berges-Gutiérrez, Fernanz-Antón, Gómez-Ríos, Ramírez-Rodríguez, García-Erce, Aldecoa.

Statistical analysis: Ripollés-Melchor, Bordonaba-Bosque, Ramírez-Rodríguez.

Obtained funding: Aseguinolaza-Pagola, Marín-Peña.

Administrative, technical, or material support:

Ripollés-Melchor, Abad-Motos, Aseguinolaza-Pagola, Padin-Barreiro, García-Orallo, Rodríguez-García, Fernanz-Antón, Ramírez-Rodríguez, García-Erce.

Supervision: Ripollés-Melchor, Abad-Motos, Díez-Remesal, Aseguinolaza-Pagola, Padin-Barreiro, Marín-Peña, Ferrando-Ortolá, Ramírez-Rodríguez, García-Erce, Aldecoa.

Conflict of Interest Disclosures:

Dr Ripollés-Melchor reported receiving personal fees from Edwards Lifesciences and Fresenius Kabi

outside the submitted work. Dr Bisbe reported receiving personal fees from Vifor Pharma outside the submitted work. Dr Abad-Gurumeta reported receiving personal fees from Merck & Co, Ferrer Pharmaceuticals, Braun PharmaCare, Panamericana Editorial, 3M Company, Rovipharm, and Altan Pharma Ltd outside the submitted work.

Dr Jiménez-López reported receiving personal fees from Edwards Lifesciences outside the submitted work. Dr Marín-Peña reported receiving personal fees from Johnson & Johnson, 3M Company, and Stryker Corporation outside the submitted work. Dr Cassinello-Ogea reported receiving personal fees from Vifor Pharma outside the submitted work. Dr García-Erce reported receiving personal fees from Alexion Pharmaceuticals, Inc, Amgen, Inc, Braun PharmaCare, Celgene Corporation, Ferrer Pharmaceuticals, GlaxoSmithKline, Inmucor, Jansen Pharmaceutica, Novartis International AG, Octapharma AG, Sanofi, Sandoz, Inc, Terumo Corporation, Vifor Pharma, and Zambon and nonfinancial support from Vifor Pharma outside the submitted work. No other disclosures were reported.

Funding/Support: This study was supported by institutional and/or departmental sources. The Postoperative Outcomes Within Enhanced Recovery After Surgery Protocol in Elective Total Hip and Knee Arthroplasty (POWER2) study was supported by the Spanish Perioperative Audit and Research Network (REDGERM).

Role of the Funder/Sponsor: The sponsors had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Group Information: All researchers of the POWER2 Study Investigators Group collected data and provided care for study patients. The POWER2 Investigators Group includes the following centers and investigators: **Hospital Universitario Nuestra señora de Candelaria, Santa Cruz de Tenerife:** María Elena Espinosa-Domínguez (principal investigator [PI]), José L. Pozo-Gómez, and María E. Yanes-Luque (data validator) (Department of Anesthesia). **Hospital Universitario de Jaén, Jaén:** Alejandro Martínez-García (PI), Almudena Morales-Rojas, Ángela M. Soriano-Pérez, Enrique García-Vena, María Navío-Poussivert, Laura Rodríguez-Melguizo, Sonia Gutiérrez-Jiménez, and Juan M. Ortiz-Cortés (data validator) (Department of Anesthesia). **Hospital Universitario del Sureste, Arganda del Rey:** Diego Anta-Redondo (PI) and Raquel Álvarez-Manzanedo (data validator) (Department of Anesthesia). **Complejo Hospitalario Universitario Insular Materno Infantil de Gran Canaria:** Dolores Betancort-Gutiérrez (PI), Abraham Ortega-Ramos, Agustín Fuentes-González, Alexis Rodríguez-León, Cristina Martín-Rico, Juan José Rodríguez-Ruiz, Laura Concepción-Santana, Rut Aracil-González, Luiz Beltrame, Patricia Borrego-Ojeda, Raquel Sánchez-Nuez, Oliver Monzón-Déniz, and José Valín-Martínez (data validator) (Department of Anesthesia). **Hospital Universitario La Paz, Hospital Cantoblanco, Madrid:** Hanna Pérez-Chrzanowska (PI) and María D. Burgueño-González (data validator) (Department of Anesthesia). **Hospital Universitario Miguel Servet, Zaragoza:** María C. Cassinello-Ogea (PI),

María C. Fraca-Cardiel, María P. Herranz-Andrés, and Inmaculada Gonzalo-Pellicer (data validator) (Department of Anesthesia). **Consorci Sanitari del Garraf:** Fernando Remartínez-Fernández (PI) and Nicolás M. Decoud-Paniagua (data validator) (Department of Anesthesia). **Agencia Sanitaria Costa del Sol, Marbella:** María P. Castillo-Mafla (PI) and Adrián Santos-Macías (data validator) (Department of Anesthesia). **Hospital García Orcyoyen, Estella, Navarra:** María Remón-Izquieta (PI and data validator), Guillermo Napal-Ongay, Gabriel Cerdán-Rodríguez, and Inés Plaja-Martí (Department of Anesthesia). **Hospital Juan Ramón Jiménez, Huelva:** Juan V. Lorente-Olazábal (PI and data validator), Elena Benito-Ramallo, José L. Bonilla-García, María D. Díaz-Lara, José M. García-García, Alejandro Gavira-Leon, María De-la-Peña-Gómez-Dominguez, Pablo Longo-Guridi, Marisol Hernández-del-Castillo, Irene Jarana-Aparicio, Irene Mojarro-Zamora, Cecilia Prieto-Candau, Ana M. Quintero-Moreno, and David Soriano-López (Department of Anesthesia). **Hospital Universitario Quirón Dexeus, Barcelona:** Sergi Alabau-Rodríguez (PI) and Ariadna Subirà-González (Department of Anesthesia); Carlomagno Cárdenas-Nylander (PI and data validator) (Department of Orthopedic Surgery). **Hospital Clínico Universitario San Juan de Alicante:** Jorge Hernando-Saez (PI), Sergio Tejada-Ortega, Paula Arnau-Papí, Francisco Manresa-Ballester, Javier Agulló-Agulló, Luis Such-Sánchez, Vera García-García, Carolina Fernández-Ortuño, and María José Guillamón-Marín (data validator) (Department of Anesthesia). **Hospital Universitario Lucus Augusti:** María Bermudez-López (PI and data validator), María López-Rodríguez, Patricia Vázquez-Dopazo, Gloria Prado-Núñez, and Carlos Villagrán-Marín (Department of Anesthesia). **Complejo Hospitalario Universitario de Ourense:** Leticia Gómez-Viana (PI and data validator), Olalla Figueiredo-González, David García-Quiroga, Eva López-López, Jorge Olmo-Do-Rodríguez, Ariadna Rodríguez-Rodríguez, Raquel Ruido-Dacal, and Eva Villar-Arcay (Department of Anesthesia); Roberto Casin-Bricio (Department of Orthopedic Surgery); Fernando Lamas-López (Department of Dental and Oral Medicine). **Complejo Hospitalario Universitario de Pontevedra:** Marina Varela-Durán (PI and data validator), Pilar Díaz-Parada, Beatriz Hernández-Vázquez, Cristina Barreiro-Pardal, Ángela González-Vilar, Marina Barreiro-Torres, and Susana Marcelo-Brage (Department of Anesthesia). **Hospital General Universitario Reina Sofía, Murcia:** Antonio Murcia-Asensio (PI), Abdalaziz Lanagran, Iván Morales-González, José Fernando-Moreno, Raquel Lax-Pérez, Marina Sánchez-Robles, Raúl Vilaverde-Sarabia, and Francisco Ferrero-Manzanal (data validator) (Department of Orthopedic Surgery). **Hospital Universitario Sierrallana, Torrelavega:** Silvia García-Orallo (PI and data validator), Carmen Martínez-Mielot, Ana Santillán-Fernández, and Milagrosa Ruiz-de-Infante (Department of Anesthesia); Melodia Molinero-Montes (Department of Orthopedic Surgery); Maialen Mozo-Segurado (Department of General Surgery). **Hospital Universitario Marqués de Valdecilla, Santander:** Ana Solar-Herrera (PI and data validator), M. Puerto Alcalde-Hierro, Cristina Salomón-Felechosa, María I. Sampedro-González, and Raquel Sánchez-De Arriba (Department of Anesthesia). **Hospital de Viladecans:** Ana

Faura-Messa (PI), Augusto Gómez-Paz (PI), Vicens Rodríguez-Bustamante, and Dani López-Ruiz (data validator) (Department of Anesthesia); Marc Ramos-Villanueva (Department of Orthopedic Surgery). **Hospital General de Villalba:** María Fernández-Rodrigo (PI and data validator), Jenny Ballestas-Pattety, Jose L. Gracia-Martínez, Jorge L. Martínez-Gómez, Carmen Mendoza-Jiménez, Juan C. Sánchez-Gil, and María E. Useche-Castañeda (Department of Anesthesia). **Complejo Hospitalario de Segovia:** Francisco J. García-Miguel (PI and data validator), Raquel Gómez-Sanz, Adriana S. Ramírez-Caldón, Beatriz Matji, and Pedro Román-Puche (Department of Anesthesia). **Hospital General de Elda:** María C. Ribera-Montes (PI), Calixto Sánchez-Pérez (data validator), Cristina Embid-Román, María V. González-Latorre, Cristina Pastor-Lain, Sergio Sánchez-Cortés, Jordi Sabater-Pavia, Aurora Rico-González, and Blanca González-Navarro (Department of Anesthesia). **Hospital FREMAP Madrid:** Gabriel Ruiz-Córdoba (PI and data validator), Isabel Cristóbal-García, Paula Martínez-Fariñas, and Antonio Crespo-García (Department of Anesthesia). **Hospital Universitario de Fuenlabrada:** Cristina Gil-Lapetra (PI), Fernando Setien-Moreno (PI), Beatriz Bolzoni-Marciel, Ana Z. Castro-Costoya, Andrea Cuadrado-Mancy, María I. Herrera-López, and Jose Olarra-Nuel (data validator) (Department of Anesthesia); Vicente Calzada-Exposito, Francisco J. García-Lázaro, and Carlos López-Orosa (Department of Orthopedic Surgery). **Hospital Universitario de Getafe:** María M Caro-Cascante (PI), Dolores Calles-Gato, Enrique Roca-Castillo, María J. Jiménez-García, Rocío Castellanos-González, Soledad Cortés-Mallavibarrera, Sonia Martín-Ventura, and Carmen García-Molina (data validator) (Department of Anesthesia); Segundo J. López-Gutierrez (PI) (Department of Orthopedic Surgery). **Althia Xarxa Assistencial Universitària de Manresa:** Carla Farré-Tebar (PI), Francesca Reguant-Corominas, Marcelo Borderas, Montserrat Cazorla-López, Belén Gil-Calvo, Isabel Pérez-Reche, and Pere Esquius-Jofré (data validator) (Department of Anesthesia). **Hospital Universitario de Basurto, Bilbao:** José C. Herrero-Herrero (PI and data validator), Manuel E. Ballesteros-de-Arco, and Naiara Pérez-Vila (Department of Anesthesia). **Hospital Bidasoa:** Carlos Yárrnoz (PI and data validator), Izaskun Emazabel (PI and data validator), María J. Asensio, Francisco J. Estomba, Itziar Murguía, and Juan R. Salmerón (Department of Anesthesia). **Hospital Clínic de Barcelona:** Nuria Martín (PI), Misericordia Basora, Fátima Salazar, Montserrat Tió, Adriana Martínez-Camacho, Carlos Ferrando-Ortolà, and Xavier Sala-Blanch (data validator) (Department of Anesthesia); Jenaro A. Fernández-Valencia-Laborde and Luis Lozano (Department of Orthopedic Surgery). **Hospital Clínico Universitario de Valladolid:** María A. Miñambres-Villar (PI and data validator), Rita P. Rodríguez-Jiménez (data validator), Sandra Fernández-Caballero, Nuria Ruiz, and Sara Cocho-Crespo (Department of Anesthesia); Ana Zabalza-Peláez and Aurelio Vega-Castillo (Department of Orthopedic Surgery). **Hospital Universitario Virgen de la Salud-Toledo:** Filadelfo Bustos-Molina (PI and data validator), Fátima Ortiz-Parra, María Martín-Carrillo, María Gimeno-García, María Jesús Martín-Jiménez, Paula Álvarez-Buitrago, Susana Gallardo-Sánchez, Itziar

Insausti-Pacheco, Marta Torres-Montalvo, and María V. Fernández-Roldán-Galán (Department of Anesthesia). **Complejo Hospitalario Universitario de Albacete:** Ana Checa-Betegón (PI), Cecilia Ais-Dávila, Pedro M. Canales-Lara, Beatriz Carretero-de-la-Encarnación, Carlos Martínez-Villar, Elena Simón-Polo, and José A. Monsalve-Naharro (data validator) (Department of Anesthesia). **Complejo Hospitalario de Mérida:** José M. Tena-Guerrero (PI and data validator), Estefanía Palma-González (PI), and Enrique González-Sánchez (Department of Anesthesia). **Complejo Hospitalario Universitario A Coruña:** Manuel Á. Gómez-Ríos (PI and data validator), Eva Mosquera-Rodríguez, Sara delRío-Regueira, Sonia González-Bardanca, and Felisa Álvarez-Refojo (Department of Anesthesia). **Hospital de Dénia Marina Salud:** Clara Dosset-Izaguirre (PI) and Gloria Coronel-Sendra (PI and data validator) (Department of Anesthesia). **Hospital General Universitario de Elche:** José L. Muñoz-Rodes (PI and data validator), Raquel Flores-Mohedano, Vicent Cuquerella-Martínez, Marta Resalt-Pereira, José M. Ronda-Pérez, and Carlos Picó-Rodríguez (Department of Anesthesia). **Hospitales del Parc de Salut Mar:** Elvira Bisbe (PI and data validator), Sandra Aguado, Luis Carol, Carlo Moreno, and Ana Pardo (Department of Anesthesia); Alex Coello, Marc Tey, and Fernando Marques (Department of Orthopedic Surgery). **Hospital Universitario de Jerez de la Frontera:** Jacobo José Fedriani-de-Matos (PI and data validator), Ana Moreno-Martín, Ana Martínez-de-Castro, Elena García-Gómez, Fátima García-Herrera, María Muñoz-Zambrano, María Pérez-García, Marta Ruiz-Mayo, and María Milagrosa Santana-Pineda (Department of Anesthesia). **Hospital Universitario de La Plana, Villareal:** Juan C. Rivera-Gaitero (PI), Belen Poyatos-Robles, Carmen Alcalde-Sánchez, José P. Calabuig-Soler, María Á. Daras-Orenga, Montiel Redondo-Castillo, Paloma Ripollés-Martínez, and Ana Nicolau-Gozalbo (data validator) (Department of Anesthesia). **Hospital de Alcañiz:** Ana Maurain-Vilorio (PI and data validator), Monserrat Alsina-Pujol, Pilar Igado-Insa, Patricia Calderón, and Manuel Yaya-Beas (Department of Orthopedic Surgery). **Hospital Central de la Cruz Roja, San José y Santa Adela:** Cristina Massa-Gómez (PI and data validator), Carmen Lope-Prado, Sonsoles Martín-Alcrudo, Julio A. Paz, Guadalupe Población-García, Luis F. Simón, and Alieriza Tabatabaian (Department of Anesthesia). **Hospital Clínico Universitario de Valencia:** Ana Mugarra-Llopis (PI and data validator), Estefanía Gracia-Ferrándiz, Esther Romero-Vargas, Ferran Serralta-Cabedo, and Ana Jurado-Pulgar (Department of Anesthesia). **Hospital Comarcal d'Ampostà:** Juan Carlos Sanz-López (PI) and Estanislao Manuel-Farrés (data validator) (Department of Anesthesia). **Hospital General Universitario de Ciudad Real:** Silvia Gil-Trujillo (PI and data validator), Remedios Moreno-Ballesteros, Juan D. Valencia-Echeverri, Juan José Arroyo, Tatiana Castellar-Cuesta-Cabezas, and Víctor Baladrón-González (Department of Anesthesia). **Hospital Universitario La Paz:** Alejandro Suarez-de-la-Rica (PI and data validator), Sofía Díaz-Carrasco, Laura Fernández-Benítez, and Almudena Martín-Martín (Department of Anesthesia). **Hospital Universitario Quirón Madrid:** Rocío Díez-Munar (PI and data validator) and África Fernández-Sánchez (Department of

Anesthesia). **Hospital Santa Clotilde de los Hermanos de San Juan de Dios:** Francisco Barrios (PI and data validator) and Manuel Buitrago-Alonso (Department of Anesthesia). **Hospital Universitario San Juan de Reus:** Lydia Cortés-Gómez (PI and data validator) and María P. Sala-Francino (Department of Anesthesia); Marta Llauredó-Pou and Alfredo Rodríguez-Gangoso (Department of Orthopedic Surgery). **Hospital Universitario de Bellvitge:** María V. López-Reig (PI and data validator), Xavier Balderas-González, Javier Bocos-García, María J. Colomina-Soler, and Guillermo Puig-Sanz (Department of Anesthesia); Marcos del-Carmen, Daniel Rodríguez-Pérez, and Carles Tramunt-Monsonet (Department of Orthopedic Surgery). **Hospital Universitario de Canarias:** Josué Nodarse-Medina (PI and data validator), María P. Rubio-Florez (PI), Fernando Hernández-Zayas, Paula Pineda-Bolivar, Ramón M. Orza-Muñiz, and Janet C. Hernández Chávez (Department of Anesthesia). **Hospital Universitario de Móstoles:** Beatriz Nacarino-Alcorta (PI and data validator), Mercedes Martínez-García (data validator) (Department of Anesthesia), Eugenio Ferrer-Santacruce (PI), and Pablo Cendrero-Cendrero (Department of Orthopedic Surgery). **Hospital Universitario Sanitas La Moraleja, Madrid:** Roberto Ruiz-Abascal (PI), Alfonso Moreno-Sánchez-Cañete, Amaya Moreno-Borque, Ana Argüelles-Baquero, Borja San-Pedro-de-Urquiza, David Salvatierra-Díaz, Silvia Ordóñez-Díaz, and Antonio López-del-Castillo (data validator) (Department of Anesthesia). **Hospital Universitario de Burgos:** Beatriz Blanco-Del-Val (PI and data validator), Sergio Bascones-Nestar, María J. Cuesta-Agudo, María I. Fernández-Martínez, María A. Gutiérrez Gómez, Ana M. Manzanal-Bañuelos, Blanca Delia-Martínez-Castro, Elena Martínez-Ibeas, María P. Isasi-Nebreda, José A Tudanca-Martínez, Juan M. de-Vicente-Lorenzo, Cristina Villalain-Pérez, and Rodrigo Sanllorente-Sebastián (Department of Anesthesia); María E. López-Díez and María I. Mora-Fernández (Department of Orthopedic Surgery). **Hospital Universitario Central de Asturias, Oviedo:** Ana B. Cuéllar-Martínez (PI and data validator), Rosa A. Álvarez-Fernández (PI), Ángela García-Díaz-Negrete, Helena García-Sánchez, and Beatriz Ayuso-Iñigo (Department of Anesthesia). **Hospital Universitario Infanta Cristina:** Fernando Pablo Somoza-Sáez (PI and data validator) (Department of Anesthesia). **Hospital Universitari Joan XXIII, Tarragona:** Ana M. Fuentes-Baena (PI and data validator), María Miranda-Marín, Ione Montalvo-Aguirrezabala, Raúl Puig-Bitria, Carlos Ros-Góngora, and Judit Saludes-Serra (Department of Anesthesia). **Hospital General de La Palma:** Johnycar Pérez-Morales (PI), Rafael A. Omaña-García (PI and data validator), Rafael Bello-Puentes, Ana P. Castillo-German, Jerzy Tadeusz-Dwojak, Mary Ursula Iribarren-Adan, Anibal Ricardo Pérez-García, Rita C. Quiroz-Cornejo, Dieter Weingaertner, and Mauricio German-Martínez (Department of Anesthesia). **OSI Alto Deba:** Pablo Renedo-Corcóstegui (PI and data validator) (Department of Anesthesia). **Hospital de Palamós:** Diana L. Mariñansky (PI), Onel Morales-Torres (PI and data validator), Eduardo Agusti-Cobos, and Silvia Sague-Bou (Department of Anesthesia). **Hospital de La Princesa, Madrid:** Jara Torrente-Pérez (PI) and Enrique Alday-Muñoz (data validator) (Department of Anesthesia); Pablo

Peñalver-Andrada (Department of Orthopedic Surgery). **Hospital Quirón Zaragoza:** Pedro Arauzo-Pérez (PI and data validator) and Paula Arauzo-Casedas (PI) (Department of Anesthesia). **Hospital Universitario Rey Juan Carlos, Móstoles:** Juan C. Estupiñán-Jiménez (PI), Eliomara J. Cabello-Urbina, and Irene Portero-Larragueta (data validator) (Department of Anesthesia). **Hospital de Sagunto:** Raúl Duplá-Pérez (PI and data validator), Rafael de-Diego-Catalá, and Marianela Martínez-Hermosilla (Department of Anesthesia). **Hospital Sant Joan Deu-Palma/Mallorca:** David Fabián-González (PI), Pedro José Benito-Rodríguez (PI), Ana Dora-Parera, Miguel Ferrá-Verdera, Nicolás García-de-Paredes-Porro, Luis M. Marco-Alós, Luis C. Mora-Fernández, Carlos Raduan-Ripoll, Cristina Sansaloni-Perelló, José V. Torres-Pérez (data validator), and Matias J. Sastre-Nadal (Department of Anesthesia). **Hospital de la Santa Creu i Sant Pau, Barcelona:** Rolf Hoffmann (PI), Marisa Moreno-Bueno, Marta Argilaga-Nogués (data validator), Adrià Font-Gual, Mireia Rodríguez-Prieto, Rafael Anaya-Camacho, Victoria Baños-Lapuente, José M. Gil-Sánchez, Marc Bausili-Ribera, Rubén Díaz-Jover, María M. Felipe-Correoso, Laura Parrilla-Quiles, Silvia Barbero-Espinosa, Xavier Crusi-Sererols, Xavier Aguilera-Roig, Màrius Valera-Peretgàs, Marcos Jordán-Sales, José C. González-Rodríguez, Esther Moya-Gómez, and Sandra Huguet-Miguélez (Department of Orthopedic Surgery). **Hospital Universitario Son Espases:** David Fabián-González (PI and data validator), Leandro Augusto-Brogí, Irene Enriquez-Valcarcel, Neus Esteve-Pérez, Germán Gómez-Romero, Carmen Melero-Larrocha, Grineldy Mercedes-Guzman, Elena Monterroso-Garrido, and Luis C. Mora-Fernández (Department of Anesthesia). **Clínica Universidad Navarra Madrid:** María M. Álvarez-Fernández (PI) and Elisa Fernández-Elias (data validator) (Department of Anesthesia). **Hospital de Villajoyosa, Alicante:** Alejandra Molines-Cantó (PI), Anna Grabowska, Cristina Muncck, Gracia Barber-Ballester, María A. Korobkoff-Fernández, María A. Lengua-Egaña, María I. Tarí-Bas (data validator), and Patricia Valls-Linares (Department of Anesthesia). **Hospital Universitario Virgen Macarena, Sevilla:** Héctor Berges-Gutiérrez (PI and data validator), Miguel Aisa-Gasca, José D. Bautista-González, Francisco J. Carballido-Pascual, Aurora Cruz-Ortega, Jorge García-García, Beatriz Gullón-Descane, Mercedes Gutiérrez-Moreno, Liena Izquierdo-López, Yolanda Lozano-Guijarro, Estefanía Peralta-Espinosa, and Ana Rodríguez-Artacho (Department of Anesthesia). **Hospital Arnau De Vilanova, Valencia:** Enrique Lloria-Pons (PI and data validator), Francisco Gramuntell, José L. Carrión, and Sara Alegre (Department of Anesthesia); Inmaculada Montón and Javier García (Department of Orthopedic Surgery). **Hospital De Liria:** Cristina Ruiz (PI and data validator), Jaime García, Juan Carrizo, and Miguel Á. Gil (Department of Anesthesia); Susana Moro and Daniel Herrero (Department of Orthopedic Surgery). **Hospital Universitario de Cruces, Bilbao:** Alberto Sánchez-Campos (PI), Xabier Ormazabal-Rementería, and Alberto Martínez-Ruiz (data validator) (Department of Anesthesia). **Hospital Universitario de Álava, Vitoria-Gasteiz:** Margarita Logroño-Ejea (PI and data validator), Alaitz Urriaga-Urrestizala, Erika Olea-De-La-Fuente, and Lucía Pérez-De-Albeniz-Vesga (Department of

Anesthesia); Alexis I. Fernández-Juan, Borja López-Hernando, Luis Á. Minguéz-Cabeza, and Jorge Guadilla-Arsuaga (Department of Orthopedic Surgery). **Hospital de Sant Joan Despi Moisés Broggi:** Jesús Fernanz-Antón (PI), Lucía Gil-Gómez, and Sandra Marmaña-Mezquita (data validator) (Department of Anesthesia); Xavi Bial-Vellvé and Carlos X. Rojas-Coutto (Department of Orthopedic Surgery); Carlos Jericó-Alba (Internal Medicine). **Hospital del Dos de Maig:** Sandra Marmaña-Mezquita (PI and data validator) (Department of Anesthesia); Borja García-Torres and Vicente J. López-Pérez (Department of Orthopedic Surgery). **Consorcio Corporació Sanitària Parc Taulí, Sabadell:** Anna Peig-Font (PI and data validator), Andrea Vallejo-Tarrat, Estefanía Rabaneda-Romero, Silvia Juncà-Martí, and Laura Ricol-Lainez (Department of Anesthesia); Albert Soler-Cano, Christian Yela-Verdú, Juan Peñalver, and Vanesa Ros-Munne (Department of Orthopedic Surgery); Helena Costa and Francisco Zamora-Carmona (Nursery). **Hospital Universitario Donostia Unibertsitate Ospitalea:** Marta Aseguinolaza-Pagola (PI and data validator), Ainhoa Aginaga-Badiola, Alazne Enparantza-Aiestaran, Rebeca Alcalde-León, Ane López-Montecelo, Anna Coll-Triquell, Belén-Barandiaran-Benito, Berta Castellano-Paulis, Ibonne Barceló-Zarraga, Elena del-Val-Peciña, María J. Goenaga-Álvarez, Manuel Eced-Sanchez, Nagore Echeverria-Rodríguez, Nuria González-Jorrín, Olga Cengotita-Blanco, and Ainhoa Garmendia-Odrizola (Department of Anesthesia). **Hospital Universitario Fundación Alcorcón:** Míriam Sánchez-Merchante (PI and data validator), Alma M. Espinosa-Moreno, Patricia Gaité-de-Vicente, Santiago García-del-Valle-y-Manzano, Livia Gisbert-de-la-Cuadra, Anna Katarzyna Grzanka, Violeta M. Heras-Hernando, Beatriz Martín-Vaquero, Alfredo Plaza-Mardomingo, Roberto Real-Valdés, and Pablo Redondo-Martínez (Department of Anesthesia). **Hospital Universitario Rio Hortega, Valladolid:** Cesar Aldecoa (PI and data validator), Irene García-Saiz, Eduardo Santos, Daniel Almudi, Clara Bolaños, and Borja Morales (Department of Anesthesia). **Hospital Universitario Ramón y Cajal, Madrid:** Yolanda Díez-Remesal (PI and data validator), Alberto José Balvis-Balvis, Alberto Berruezo-Camacho, Inés de-la Hoz-Polo, Borja Hinojal-Olmedillo, Sara Martín-Burcio, María C. Martín-González, and Marta Ramírez-González (Department of Anesthesia). **Hospital Ernest Lluch Calatayud:** Leticia I. Pérez-Pascual (PI and data validator) and Marta Ortiz-Enciso (Department of Anesthesia); Alfredo Á. Rodríguez-León and María T. Domínguez-Ayasa (Department of Orthopedic Surgery). **Hospital Universitario Dr Peset, Valencia:** Jorge Bellver (PI), Joana Baldó, Cristina Álvarez, Concepción Barrachina, José M. Castillo (data validator), José Higuera, Estefanía Martínez-González, María J. Molina, Leticia Narváez, Oscar Ramírez, Fernando Rauer, María T. Richart, Amparo S. de-Merás, Cynthia Sánchez-Martín, and Luisa Sapena (Department of Anesthesia). **Hospital Universitario de Cáceres:** Juan M. Redondo-Enriquez (PI and data validator), Francisco B. Amador-Penco, Antonio Barba-Orejudo, Inés de-Tena-Rocha, Marta Donoso-Domínguez, Laura Fuentes-Coco, Fernando García-Montoto-Pérez, Yasmina González-Marín, Luis G. Smith-Vielma, Álvaro San-Antonio-Gil, Pedro Fernández-Nuñez, and José

I. Hermoso-Martínez (Department of Anesthesia). **Hospital Universitario De Cabañes:** Javier Albaladejo-Magdalena (PI and data validator), Ana Rodríguez-Morala and Javier Carbayo-Lázaro (Department of Anesthesia). **Clínica Santa Cristina, Albacete:** Ubaldo Vicente-Gutiérrez (PI and data validator) (Department of Anesthesia). Jesús Martínez-Casrtoverde and Eduardo Rodríguez-Vellando-Rubio (Department of Orthopedic Surgery). **Clínica HLA Vistahermosa:** Jesús J. Mas Martínez (PI and data validator) and Javier Sanz-Reig (PI) (Department of Orthopedic Surgery); Enrique Cardenas-Alegría (Department of Anesthesia). **Complejo Asistencial De Zamora:** Ana M. García-Sánchez (PI and data validator), María Gómez-Fernández, Ana M. Lafuente-Lera, and Gema Martínez-Ragüés (PI) (Department of Anesthesia); Gonzalo Humada-Álvarez and Camino Román-Dopazo (Department of Orthopedic Surgery). **Complejo Hospitalario Universitario de Badajoz:** María D. Torrado-Criado (PI and data validator), Laura Caperote-Anchez, Juan R. Caro-González, Francisco S. Castaño-García, Ana Marín-Moreno, Gema Montero-Mejías, and Cristina Pinilla-Silva (Department of Anesthesia); Rafael Lorente-Moreno (data validator) (Department of Orthopedic Surgery). **Complejo Hospitalario Universitario De Vigo-Eoxi Vigo:** Lidia Padin-Barreiro (PI and data validator), María I. Vidal Lopo, Iván Areán-González, María A. Gil-Campelo, Pedro Martínez-Gutián, José R. Reparaz-Andrade, and Gerardo Baños-Rodríguez (Department of Anesthesia). **Hospital De La Cruz Roja, Gijón:** Nicolás Rodríguez-García (PI and data validator) (Department of Anesthesia); Daniel García-Rodríguez (PI) (Department of Orthopedic Surgery). **Hospital Universitario de Gran Canaria Doctor Negrín:** Ángel Becerra-Bolaños (PI and data validator), Aurelio Rodríguez-Pérez, Iván M. Beirutti-Ruiz, Carmen Arachelly-Focaccio-Tejada, Julio Gómez-Marrero, Leonardo Cabrera-Ramírez, and Mutaz Mourad-Mohamed (Department of Anesthesia). **Hospital Don Benito Vilanueva:** Marta López-Marqués (PI), María P. Rodríguez-Chaparro, Enrique Del-Cojo-Peces (data validator) (Department of Anesthesia). **Hospital El Bierzo:** Sergio Núñez-Moreno (PI), Sarai Suárez-Fernández, and Irene Ruperez-San-Emeterio (data validator) (Department of Anesthesia); Marta Novoa-Cortiñas and Alicia Sánchez-Parada (Department of Orthopedic Surgery). **Hospital De Galdakao-Usansolo:** Pilar Pérez-Sánchez (PI and data validator), Claudia Velasco-Oficialdegui, Fernando Jáuregui-Beraza, Pilar Macua-Biurrun, Raquel Malanda-Callejo, and Aizpea Aguirrezabal-Juaristi (Department of Anesthesia). **Hospital General de Granollers:** Germán Moreno-Aguilar (PI and data validator), Carmen Sanabria-Pico, and Regina Sopena-García (Department of Anesthesia). **Hospital General Universitario Gregorio Marañón, Madrid:** Roberto de-Diego-Fernández (PI), Alba Burgos-Santamaría, Carmen Corbella-Jiménez (data validator), Elena de-la-Fuente-Tornero, Elena González-Benito, Fernando Higuerro-Cantenero, David Martínez-Gascuña, Ángeles de-Miguel-Guijarro, Ana G. Pizarro-Calderón, Mercedes Power-Esteban, Usamah Rabih-Kanji, Olalla Varela-García, Juan M. de-Vicente-Peña, and Santiago García-Hernández (Department of Anesthesia). **Hospital Universitario de Guadalajara:** Mercedes Cabellos-Olivares (PI), Laura M. Martínez-Meco (PI),

and José R. Rodríguez-Fraile (data validator) (Department of Anesthesia). **Hospital Universitario 12 de Octubre, Madrid:** Álvaro Ramiro-Ruiz (PI and data validator), Pablo García-Pimentel, Sergio García-Pérez, Noelia González-Rabadán, Adrián Martínez-López, Teresa Martínez-Marivela, Belinda Montalbán-Moreno, Noelia Zurera-Plaza, Rubén Salas-Rodríguez, Claudia Sarraís-Polo, and Claudia Olea-Vielba (Department of Anesthesia). **Hospital d'Igualada:** Laura Perelló-i-Cerdà (PI), Josep M. Bausili-Pons (data validator), Ana B. Pedregosa-Sanz, Ana M. Pedrero, and Joaquim Sabater-Recolons (Department of Anesthesia). **Hospital Universitario Infanta Leonor, Madrid:** Javier Ripollés-Melchor (PI), Ane Abad-Motos (PI), Beatriz Nozal-Mateo, Rut Salvachua-Fernández, Alfredo Abad-Gurumeta, Elena Sáez-Ruiz, Eugenio Martínez-Hurtado (data validator), Elena Lucena-de-Pablo, Elena Nieto-Moreno, and Norma Aracil-Escoda (Department of Anesthesia); Oliver Marín-Peña (PI) and David Murillo-Vizuet (Department of Orthopedic Surgery); Elena Sánchez-Pérez (Nursery). **Hospital General Universitario de Castellón:** José M. España-Pamplona (PI and data validator), Laura Jordá-Sanz, María Vellida-Vilarroig, and María I. De-Miguel-Cabrera (Department of Anesthesia). **Hospital Universitario Clínico San Carlos, Madrid:** Rubén Sánchez-Martín (PI and data validator), Jesús Hurtado-Martín, Jhon López-Hormiga, Julián A. Vargas-Ardila, Karina Martínez-Cruz, Lourdes Durán-Giménez-Rico, and Rosalía Navarro-Pérez (Department of Anesthesia). **Fundación Salud Emporda - Hospital De Figueres:** Manel Roig-Gasull (PI and data validator), María del Mar Piazuelo, and Carlos Salazar-Zamorano (Department of Anesthesia). **Consorcio Hospital General Universitario de Valencia:** Juan C. Catalá-Bauset (PI), Rafael Colomina-Rodríguez (data validator), Lorenzo Hernández-Ferrando, Pablo Rodríguez-Gimillo, Raquel Moratalla-García, Elena Biosca-Pérez, María Olivas-Dols, Lucas Rovira-Soriano, Rubén Rubio-Haro, Francisco Gil-Chaves, Francisco Jara-García, and Jorge Ubeda-Pascual (Department of Anesthesia). **Hospital Manises:** Aurora Moreno-Gazquez (PI), Raquel Viñedo-Lillo, Óscar Torres-Rico, and Vicente R. Chisbert-Cuenca (Department of Anesthesia). **Hospital Municipal de Badalona:** Eduard Siles-Fuentes (PI), Ester Mora-Solá, and Javier González-López (data validator) (Department of Orthopedic Surgery). **Hospital Quirón de Marbella y Hospital Quirón Campo de Gibraltar:** Blas Rodríguez-Gil (PI and data validator), Belén Castillo-Cardona, Raimundo Escudero-Concha, Mathieu Juvin-Bouvier (PI), and José Romero-Aguilar (Department of Anesthesia). **Hospital Quirónsalud Torreveja:** Andrej Balik (PI and data validator), Noelia Ruiz-Vera, Julio P. Vinuesa-Núñez, and Ana B. Tovar-López (Department of Anesthesia). **Hospital Universitario San Agustín, Avilés:** Diana de-la-Uz-Díaz (PI), Paz Alonso-Pérez (data validator), Juan R. Artamendi-Suárez, Jesús Bujarrabal-Martínez, Lorena España-Fuente, Ángel Gómez-Martín, María J. González-Miranda, Covadonga Gutiérrez-González, Beatriz Laserna-Cocina, Marta López-Pérez, Andrea Manzano-Patallo, Gabriel Mella-Pérez, Rosa E. Méndez-Redondo, Marina Rodríguez-Rodríguez, Cristina M. Rodríguez-Rodríguez, Laura Morante-Carrión, and Pablo de-la-Rica-Fernández

(Department of Anesthesia). **Hospital Universitario Mútua Terrassa:** Marga Novellas-Canosa (PI) (Department of Anesthesia); Francesc Anglès-Crespo (PI and data validator) (Department of Orthopedic Surgery). **Hospital Universitario Severo Ochoa, Leganés:** Vanesa Blázquez-Calvo (PI), Luis Fuentetaja-Martín-Portugués, Patricia Lloreda-Herradón, Adela Gardeta-Pallarés, Soraya Gholamian-Ovejero, Raquel González-Velasco, Adolfo Ramos-Luengo, Marta Vicente-Orgaz, Purificación Sánchez-Zamora, and Gema Fraga-Casais (data validator) (Department of Anesthesia); Hilario Miguez-Pérez (Department of Orthopedic Surgery). **Hospital Universitari i Politècnico la Fe de Valencia:** Lucía Cabezudo-de-la-Muela (PI and data validator), Beatriz Castel-González, Daniel Martínez-Catalá, and María D. Ruiz-Boldua (Department of Anesthesia). **Hospital Clínico Universitario Lozano Blesa de Zaragoza:** Aurora Callau-Calvo (PI), Patricia García-Consuegra, and Javier Longás-Vailén (data validator) (Department of Anesthesia); José M. Ramírez-Rodríguez (Department of General Surgery). **Hospital de Mataró:** Marta Villanova-Baraza (PI and data validator), Roser Torruella-Turró, Raquel Algilaga-Segovia, Isabel Fábregas-Blanco, María Farolera-Bonamusa, Elisa Reñé-de-Antonio, Alexia Nebot-Galindo, and Cassandra Gimeno-Grauwinkel (Department of Anesthesia). **Hospital Medina del Campo:** María T. Fernández-Martín (PI and data validator), Juan C. Álvarez-López, Beatriz Cano-Hernández, and Christian Bravo-Ovadia (Department of Anesthesia). **Hospital Universitario de Torrejón:** Elena García-Fernández (PI and data validator), Sergio Llorio-Palomino, Jesús Recio-Pérez, and José M. Requena (Department of Anesthesia). **Hospital Nuestra Señora del Prado, Talavera de la Reina:** Pablo Gimeno-Fernández (PI and data validator), José L. Gómez-Agraz, Carla Iglesias-Morales, Patricia Ordoñez-Díaz, Hilda L. Quispe-Melendez, Josely Rodríguez-Pinto, Patricia Sabin-Gutiérrez, Yolanda Sarabia-Mora, and Elzbieta Stasiak-Pikula (Department of Anesthesia). **Hospital Universitario Puerta de Hierro Majadahonda:** Lucía Ormaechea-Bolado (PI and data validator), Guillermo Arespachaga-Ramos, Marina Carmona-Prieto, Fernando Herva-Pintor, Ana Lacaba-Mazarío, and Javier Martínez-Caridad (Department of Anesthesia); Teresa Beteta-Robles, Beatriz Fernández-Maza, Marta Isabel Sanz-Pérez, and Marta Yáñez-Hernández (Department of Orthopedic Surgery). **Quiron Teknon Barcelona:** Adriana Bataller-Bassols (PI and data validator), Milen Bonev-Bonev, Amparo Gómez-Mantecon, Karen Salazar-Loaiza, and Josep Rodiera-Olivé (Department of Anesthesia); Diego Collado-Gastalver, Rafael González-Adrio, Ferran Montserrat-Ramon, David Roca-Romalde, and Miquel Sales-Pérez (Department of Orthopedic Surgery). **Hospital Ruber Internacional, Madrid:** Uxío García-Aldao (PI), Judit Benitez, Pilar Bernad, Álvaro Elícegui, Daciano Gaona, Isabel Gómez, Álvaro de-la-Vega, Juan P. Von-Walter, and Javier de-Benito (data validator) (Department of Anesthesia). **Hospital Universitario de Santa Maria, Lleida:** Jordi Colomina-Morales (PI and data validator), Isabel Parada-Avendaño (Department of Orthopedic Surgery); Reis Drudis-Morell (PI) and Montserrat Torra-Riera (Department of Anesthesia). **Hospital de Zumarraga, OSI Goierri Alto Urola:** Sandra L. Muñoz-Pérez (PI and data validator), Iñaki Roa-Martínez, Angel Beriain-Arcelus, Maitane

Rubio-de-Jesús, and Marco Amambal-Altamirano (Department of Anesthesia). **Hospital Universitario Lucus Augusti:** María Bermúdez-López (PI and data validator), María López-Rodríguez, Carlos Villagrán-Marín, Gloria Prado-Núñez, and Patricia Vázquez-Dopazo (Department of Anesthesia). **Complejo Asistencial Universitario de León:** Cristina García-Pérez (PI and data validator), Sergio Marcos-Contreras, Javier Ferrero-de-Paz, and Diana Fernández-García (Department of Anesthesia). **Servicio Navarro de Salud-Osasunbidea, Pamplona, Navarra:** José A. García-Erce (PI) (Blood and Tissue Bank). **Hospital MAZ Zaragoza:** Clara Marín-Zaldívar (PI and data validator) and Amagoia Fernández-de-Gamara-Goiricelaya (Department of Anesthesia); Julian C Segura-Mata (PI), Juan F Blanco-Gómez, Carlos M Enseñat-Schulte, Silvia Gomez-Gomez, María Llorens-Eizaguerri, Lucia Lopez-Sagasta, and Jesus Rivas-Felice (Department of Orthopedic Surgery). **Hospital Universitario de Vic:** Jordi Serrat-Puyol (PI and data validator), Teresa Planella-Rius, and Julio Roza-Baglietto (Department of Anesthesia); Laia Boadas-Gironés and Daniel Estevez-Sabogal (Department of Orthopedic Surgery). **Hospital Universitario Fundación Jiménez Díaz, Madrid:** Patricio J Amaro-Soto, Philipp Peter-Granacher, Raquel García-Ortega, Raquel Iglesias-Guitian, Fernando López-Arias, and Soledad Bellas-Cotan (PI and data validator). **Hospital Virgen de los Lirios, Alcoy:** José Luis Jover-Pinillos (PI), Vicente Climent-Peris, and José Pedro García-Belmonte (data validator). **Hospital Universitario Príncipe de Asturias, Alcalá de Henares:** Carmen Demartini-Foj (PI and data validator), Fernando Poveda-Sáenz, and Francisco Barrios (Department of Anesthesia). **Hospital Universitario Infanta Elena, Valdemoro:** Rubén Casans-Frances (PI and data validator), Thais Martínez-Mejía, Enrique Golderos, and Elena Del-Hierro (Department of Anesthesia); Francisco J. Cervero, Elena Galvez, Elena Villarrubia, Rodrigo Anton, Pablo Vadillo, Daniel López-Dorado, Raquel Serrano, Rocío Mascaraque, and Hugo Garlito (Department of Orthopedic Surgery). **Hospital del Poniente, El Ejido:** Miguel Martínez-Espinosa (PI and data validator) and Alberto Albert-Ullibarrí (Department of Orthopedic Surgery). **Hospital Universitario Virgen del Rocío:** Juan Sánchez-Peña, María de los Ángeles Luengo-Pastor, Manuel Sariago-Ortiz, Jesús Fernández-López, Juan Miguel Pérez-de-Tudela, Juan Francisco Gómez-Curiel, Manuel Agustín Castro-Marín, Alvaro Sosa-Orts, Francisco Manzano-Capitán, Rosana Guerrero-Domínguez, Francisco Bullón-Barrera, Inmaculada Jimeno-Puche, Francisco Polonio-Enríquez, Lucia Romero-Escalante, Ana Martínez-Saniger, Concepción Martínez-Torres, and Ignacio Jiménez-López (PI and data validator) (Department of Anesthesia).

Additional Contributions: Julián Álvarez Escudero, MD, PhD, President of the Spanish Society of Anesthesiology and Critical Care (SEDAR) provided a noncompensated prerecruitment announcement of the study on the SEDAR website. Alejandro Bona Enguita, BS, served as the POWER2 website webmaster. No financial compensation was provided.

REFERENCES

- Weiser TG, Regenbogen SE, Thompson KD, et al. An estimation of the global volume of surgery: a modelling strategy based on available data. *Lancet*. 2008;372(9633):139-144. doi:10.1016/S0140-6736(08)60878-8
- Kurtz SM, Lau E, Ong K, Zhao K, Kelly M, Bozic KJ. Future young patient demand for primary and revision joint replacement: national projections from 2010 to 2030. *Clin Orthop Relat Res*. 2009;467(10):2606-2612. doi:10.1007/s11999-009-0834-6
- Global patient outcomes after elective surgery: prospective cohort study in 27 low-, middle- and high-income countries. *Br J Anaesth*. 2017;119(3):553. doi:10.1093/bja/aew472
- Bozic KJ, Grosso LM, Lin Z, et al. Variation in hospital-level risk-standardized complication rates following elective primary total hip and knee arthroplasty. *J Bone Joint Surg Am*. 2014;96(8):640-647. doi:10.2106/JBJS.L.01639
- Maradit Kremers H, Larson DR, Crowson CS, et al. Prevalence of total hip and knee replacement in the United States. *J Bone Joint Surg Am*. 2015;97(17):1386-1397. doi:10.2106/JBJS.N.01141
- Jørgensen CC, Kehlet H; Lundbeck Foundation Centre for Fast-track Hip and Knee Replacement Collaborative Group. Role of patient characteristics for fast-track hip and knee arthroplasty. *Br J Anaesth*. 2013;110(6):972-980. doi:10.1093/bja/ae505
- Kehlet H. Multimodal approach to control postoperative pathophysiology and rehabilitation. *Br J Anaesth*. 1997;78(5):606-617. doi:10.1093/bja/78.5.606
- Ljungqvist O, Scott M, Fearon KC. Enhanced Recovery After Surgery: a review. *JAMA Surg*. 2017;152(3):292-298. doi:10.1001/jamasurg.2016.4952
- Antrobus JD, Bryson GL. Enhanced recovery for arthroplasty: good for the patient or good for the hospital? *Can J Anaesth*. 2011;58(10):891-894, 894-896. doi:10.1007/s12630-011-9564-9
- Raphael M, Jaeger M, van Vlymen J. Easily adoptable total joint arthroplasty program allows discharge home in two days. *Can J Anaesth*. 2011;58(10):902-910. doi:10.1007/s12630-011-9565-8
- Reilly KA, Beard DJ, Barker KL, Dodd CAF, Price AJ, Murray DW. Efficacy of an accelerated recovery protocol for Oxford unicompartmental knee arthroplasty: a randomised controlled trial. *Knee*. 2005;12(5):351-357. doi:10.1016/j.knee.2005.01.002
- Khan SK, Malviya A, Muller SD, et al. Reduced short-term complications and mortality following Enhanced Recovery primary hip and knee arthroplasty: results from 6000 consecutive procedures. *Acta Orthop*. 2014;85(1):26-31. doi:10.3109/17453674.2013.874925
- Glassou EN, Pedersen AB, Hansen TB. Risk of re-admission, reoperation, and mortality within 90 days of total hip and knee arthroplasty in fast-track departments in Denmark from 2005 to 2011. *Acta Orthop*. 2014;85(5):493-500. doi:10.3109/17453674.2014.942586
- Wainwright TW, Gill M, McDonald DA, et al. Consensus statement for perioperative care in total hip replacement and total knee replacement surgery: Enhanced Recovery After Surgery (ERAS[®]) Society recommendations. *Acta Orthop*. 2019;115(6):1-17. doi:10.1080/17453674.2019.1683790
- Soffin EM, YaDeau JT. Enhanced recovery after surgery for primary hip and knee arthroplasty: a review of the evidence. *Br J Anaesth*. 2016;117(suppl 3):iii62-iii72. doi:10.1093/bja/aew362
- Sutton E, Herbert G, Burden S, et al. Using the Normalization Process Theory to qualitatively explore sense-making in implementation of the Enhanced Recovery After Surgery programme: "it's not rocket science". *PLoS One*. 2018;13(4):e0195890. doi:10.1371/journal.pone.0195890
- Ripollés-Melchor J, Ramírez-Rodríguez JM, Casans-Francés R, et al; POWER Study Investigators Group for the Spanish Perioperative Audit and Research Network (REDGERM). Association between use of Enhanced Recovery After Surgery protocol and postoperative complications in colorectal surgery: the Postoperative Outcomes Within Enhanced Recovery After Surgery Protocol (POWER) Study. *JAMA Surg*. 2019;154(8):725-736. doi:10.1001/jamasurg.2019.0995
- von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE Initiative. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ*. 2007;335(7624):806-808. doi:10.1136/bmj.39335.541782.AD
- Elias KM, Stone AB, McGinagle K, et al; ERAS[®] Society and ERAS[®] USA. The Reporting on ERAS Compliance, Outcomes, and Elements Research (RECOVER) Checklist: a joint statement by the ERAS[®] and ERAS[®] USA Societies. *World J Surg*. 2019;43(1):1-8. doi:10.1007/s00268-018-4753-0
- Postoperative outcomes within an enhanced recovery after surgery protocol (POWER2). ClinicalTrials.gov identifier: NCT03570944. <https://clinicaltrials.gov/ct2/show/NCT03570944>. Updated March 6, 2019. Accessed X.
- Ripollés-Melchor J, Abad-Motos A, Logroño-Egea M, et al. Postoperative outcomes within Enhanced Recovery After Surgery protocol in elective total hip and knee arthroplasty: POWER2 Study: study protocol for a prospective, multicentre, observational cohort study. *Turk J Anaesthesiol Reanim*. 2019;47(3):179-186. doi:10.5152/TJAR.2019.87523
- Castor Electronic Data Capture. <https://castoredc.com>. Published 2019. Accessed September 15, 2019.
- Jammer I, Wickboldt N, Sander M, et al; European Society of Anaesthesiology (ESA) and the European Society of Intensive Care Medicine (ESICM); European Society of Anaesthesiology; European Society of Intensive Care Medicine. Standards for definitions and use of outcome measures for clinical effectiveness research in perioperative medicine: European Perioperative Clinical Outcome (EPCO) definitions: a statement from the ESA-ESICM joint taskforce on perioperative outcome measures. *Eur J Anaesthesiol*. 2015;32(2):88-105. doi:10.1097/EJA.000000000000118
- Healy WL, Iorio R, Clair AJ, Pellegrini VD, Della Valle CJ, Berend KR. Complications of total hip arthroplasty: standardized list, definitions, and stratification developed by the Hip Society. *Clin Orthop Relat Res*. 2016;474(2):357-364. doi:10.1007/s11999-015-4341-7
- Healy WL, Della Valle CJ, Iorio R, et al. Complications of total knee arthroplasty: standardized list and definitions of the Knee Society. *Clin Orthop Relat Res*. 2013;471(1):215-220. doi:10.1007/s11999-012-2489-y

26. Mehran R, Rao SV, Bhatt DL, et al. Standardized bleeding definitions for cardiovascular clinical trials: a consensus report from the Bleeding Academic Research Consortium. *Circulation*. 2011;123(23):2736-2747. doi:10.1161/CIRCULATIONAHA.110.009449
27. Deng Q-F, Gu H-Y, Peng W-Y, et al. Impact of enhanced recovery after surgery on postoperative recovery after joint arthroplasty: results from a systematic review and meta-analysis. *Postgrad Med J*. 2018;94(1118):678-693. doi:10.1136/postgradmedj-2018-136166
28. Husted H. Fast-track hip and knee arthroplasty: clinical and organizational aspects. *Acta Orthop Suppl*. 2012;83(346):1-39. doi:10.3109/17453674.2012.700593
29. Paton F, Chambers D, Wilson P, et al. Effectiveness and implementation of enhanced recovery after surgery programmes: a rapid evidence synthesis. *BMJ Open*. 2014;4(7):e005015-e005015. doi:10.1136/bmjopen-2014-005015
30. Wainwright TW, Kehlet H. Fast-track hip and knee arthroplasty: have we reached the goal? *Acta Orthop*. 2019;90(1):3-5. doi:10.1080/17453674.2018.1550708
31. Whitcomb WF, Lagu T, Krushell RJ, et al. Experience with designing and implementing a bundled payment program for total hip replacement. *Jt Comm J Qual Patient Saf*. 2015;41(9):406-413. doi:10.1016/S1553-7250(15)41052-9
32. Muñoz M, Acheson AG, Auerbach M, et al. International consensus statement on the peri-operative management of anaemia and iron deficiency. *Anaesthesia*. 2017;72(2):233-247. doi:10.1111/anae.13773
33. Jørgensen CC, Petersen MA, Kehlet H; Lundbeck Foundation Centre for Fast-Track Hip and Knee Replacement Collaborative Group. Preoperative prediction of potentially preventable morbidity after fast-track hip and knee arthroplasty: a detailed descriptive cohort study. *BMJ Open*. 2016;6(1):e009813. doi:10.1136/bmjopen-2015-009813
34. Petersen PB, Kehlet H, Jørgensen CC; Lundbeck Foundation Center for Fast-track Hip and Knee Replacement Collaborative Group. Myocardial infarction following fast-track total hip and knee arthroplasty-incidence, time course, and risk factors: a prospective cohort study of 24,862 procedures. *Acta Orthop*. 2018;89(6):603-609. doi:10.1080/17453674.2018.1517487
35. Yang Z-G, Chen W-P, Wu L-D. Effectiveness and safety of tranexamic acid in reducing blood loss in total knee arthroplasty: a meta-analysis. *J Bone Joint Surg Am*. 2012;94(13):1153-1159. doi:10.2106/JBJS.K.00873
36. Mueller MM, Van Remoortel H, Meybohm P, et al; ICC PBM Frankfurt 2018 Group. Patient blood management: recommendations from the 2018 Frankfurt Consensus Conference. *JAMA*. 2019;321(10):983-997. doi:10.1001/jama.2019.0554
37. Guerra ML, Singh PJ, Taylor NF. Early mobilization of patients who have had a hip or knee joint replacement reduces length of stay in hospital: a systematic review. *Clin Rehabil*. 2015;29(9):844-854. doi:10.1177/0269215514558641
38. Chua MJ, Hart AJ, Mittal R, Harris IA, Xuan W, Naylor JM. Early mobilisation after total hip or knee arthroplasty: a multicentre prospective observational study. *PLoS One*. 2017;12(6):e0179820. doi:10.1371/journal.pone.0179820
39. Tayrose G, Newman D, Slover J, Jaffe F, Hunter T, Bosco J III. Rapid mobilization decreases length-of-stay in joint replacement patients. *Bull Hosp Jt Dis (2013)*. 2013;71(3):222-226.